

Part II

Can there be a (stable) Small C Multielectron State of The Newpde?

Yes, It is **Composite 3e, 2P_{3/2} at r=r_H**

Review of partI

Appendix: Where does this 'Newpde' come from?

My concept: Occam's razor optimized observable (so with observer):

$$1=1X1$$

Definition of 'observability'

zz=z (eq1) algebraically defines **1,0**: the simplest (Occam's razor) assumption. So my concept is **postulate 1** (observable) z=zz with observer (\equiv noise C optimized so $\delta C=0$) added in $z'=z'z'+C$ (eq1) thus with all z' unknown except postulated $zz=z$ (**=1,0**). So

$z=0=z_o=z'$ in the iteration of **eq.1** with $\delta C=0$ gives the 2D Mandelbrot set C so a δz (sect.1).

$z=1$ thereby implies $z'=1+\delta z$ in **eq.1** with $\delta C=0$ gets 2D Dirac eq for e,v ($N \geq 1$ **observer** eq11) These $z=0$, $z=1$ steps together imply the 2D+2D= 4D **Newpde** with $\delta z=r_H=(e^2/m)10^{40N}=C$ in $\kappa_{oo}=1-r_H/r=1/\kappa_{rr}$ with large m so small C in eq1. Note eq.11 **observables** requires Newpde $\delta z'=\psi$

So all we did here is define observability (in above "my concept")

Furthermore $N=0$ **postulate 1** can also be used in a list-define math to get the real number algebra (without all those many Rel#math axioms). Eg., $1 \cup 1 \equiv 1+1$ (B2,Ch.2).

So we get both the physics (ref5) AND (rel#)mathematics from ONE postulate1, everything! We figured it out: Just **Postulate1**

Small C' boost in composite 3e is the subject of partII

That is the subject of partII.

Small C stable Newpde State 2P_{3/2} at r=r_H: Composite 3e

Table of Contents

Ch.7 Small C stable state of New pde is Composite 3e at r=r_H 2P_{3/2} h/e flux quantization z=0

Excited state Small C Paschen Back ortho (s,c,b) and para (t) energy levels

Ch.8,9 Frobenius series solution r perturbation of each individual Paschen Back energy level ortho, para (s,c,b; t) getting the particle multiplets at each level

Ch.10,11 New pde high energy Cross-sections and Nuclear Binding energies

Ch.12 Comparison and contrast of 2P_{3/2} at r=r_H with mainstream toy model gauge theories

Newpde Composite **3e** 2P_{3/2} at r=r_H Model: small C realized

Note that we derived γ, Z_0, W^+, W^- in Appendix A Of Part I So we derived the E and B fields.

7.3 Eq.11 B Field Flux Quantization In This Enclosed Current Loop

Note if a charged particle moves in loop in a field free region that surrounds another region, there is trapped magnetic flux ϕ in that region. Also we can include minimal interaction E&M momentum/ $\hbar = k \rightarrow k + eA/\hbar = eBr/\hbar$ for uniform B field. If ψ phase is a unique function on the loop then phase $kr = (eBr/\hbar)r = (eBr/\hbar)r = e(Barea)/\hbar = e\Phi/\hbar = n2\pi$. $\Phi = 4.13 \times 10^{-15}$ for integer spin. Then upon completing a closed loop the particle's wave function will acquire an additional phase factor $\exp\left(\frac{ie\Phi}{\hbar}\right)$. But the wave function must be single valued at any point in space. This can be accomplished if the magnetic flux Φ is quantized: $e\Phi/\hbar = \pi n$, $n=0 \pm 1, \pm 2, \pm 3$, so $\Phi_o = h/(2e)$. From

NIST: $2.067833848 \times 10^{-15} \text{Wb} = \Phi_0$. half integer spin $1/2$. Integer spin $2\Phi_0 = 2(h/2e) = h/e$ for the two positrons.

7.4 Ultrarelativistic Rotator.

Side View $z=0$.

From the side of the rotator the Fitzgerald contraction goes as: $r'_H = r_H \int_{\pi/2}^0 \sqrt{1 - \frac{v^2}{c^2}} \sin\theta d\theta$.

$$= r_H \int_{\pi/2}^0 \sqrt{1 - \frac{c^2 \sin^2\theta}{c^2}} \sin\theta d\theta =$$

$$= r_H \int_{\pi/2}^0 \cos\theta \sin\theta d\theta = r_H \int_0^1 x dx = r_H \frac{x^2}{2} \Big|_0^1 = r_H/2 = 2.8 \times 10^{-15} \text{m} \equiv r'_H \quad (7.1)$$

for the ortho $2P$ state observer (i.e., $2P_{3/2}$, $2P_{1/2}$) in the horizontal plane and $r_H/2 = r''$. We must repeat this integration on the end para states, the radius is shrunk by $\tau + 2(\varepsilon + \Delta\varepsilon)$ and so is nearly a point source $S_{1/2}$ state (for the observer above the circle as for the deuterium central electron sect.10.7). We next show that the jump from ortho to para must then correspond to the jump from ε to τ fractal quantum state given τ is separable and so a orthogonal state transition.

$2r_H = 2(2.81 \times 10^{-15} = 2e^2/(m_e c^2))$, **Side view** $1/2(2r_H) = r_H$,

Radius Of Proton: Average both the \cos^2 edge ($(1/2)r$) and axial width (near zero) $(2.81713/2 + 0)/\sqrt{2} = .98F = r_p$. But the spectroscopic and scattering methods that are used to measure r_p do not assume a singularity physics at $2.8F/2 = r_{Hp}/2$ so allow for the slightly larger radius in practice proton radius = $r_p = 1.1F$.

Top view $2r_H, z=1$

So only New pde e ($z=0$ $\xi_0 = \Delta\varepsilon = m_e$) is stable. The only way to get multi e particle *stable* large ξ is with the Newpde **composite** $3e$ $2P_{3/2}$ at $r=r_H$ state. That is because we have *stability* ($dt'^2 = (1 - r_H/r)dt^2$) clocks stop at $r=r_H$. That 3rd mass also reverses the pair annihilation virtual pair creation inside the r_H volume given $\sigma = \pi r_H^2 \approx (1/20)\text{barn}$ making it merely a virtual creation annihilation event. So our $2P_{3/2}$ composite $3e$ (proton) at $r=r_H$ is the *only* stable multi e composite.

Magnetic Flux Quantization For Current Around Loop

Our Newpde II \rightarrow III quadrant eq.12 rotations (appendix A4) gave us Maxwell's equations and E&M so we can apply B fields here. We also derived quantum mechanics from that Circle equation (giving eq.11). Thus we can have quantization of the B field flux $\oint \vec{B} \bullet \vec{dA} = \Phi_0 N$ Just above (and below) the coil plane toward the edge of the coil the B direction changes and the magnitude of B goes up. So some $\vec{B} \bullet \vec{dA} = \Delta\Phi$ minimum deviation from BdA for some constant $|\vec{dA}|$ above the coil plane. Given B is perpendicular to dA at the center and the radius r_H of the coil cancels out (eq.2 below) this $\Delta\Phi$ flux could be over the center where the relevant γ is needed. Thus we must write $\Sigma\Delta\Phi = \Phi = BA = B\pi r_H^2$ with the B at the center of the coil for $z=0$ (appendix). So effective r_H slightly bigger (making B smaller) but r_H cancels anyway. So

$$BA = \left(\frac{\mu_0 i}{2r_H}\right) \pi r_H^2 = \Phi_0 (\#2P_{3/2} \text{lobes}). \quad (1)$$

Also $r_H = e^2/m_e c^2$, $q/t = i$. $q = e = 1.6 \times 10^{-19} \text{C}$, $\Phi_0 = \text{NIST: } 2.067833848 \times 10^{-15} \text{Wb}$, $1/\gamma$ dilation of r_H but it and r_H get canceled out here. The time t dilation γ remains in the current 'i' moving frame of reference. Recall that for circular motion: $c = D/t = 2\pi r_H/t$ so:

$$t = \frac{2\pi r_H}{\gamma c}, \text{ so } i = \frac{e}{\left(\frac{2\pi r_H}{\gamma c}\right)}$$

$$BA = \frac{\mu_0 i}{2r_H} (\pi r_H^2) = \frac{\mu_0}{2r_H} \left(\frac{e}{\left(\frac{2\pi r_H}{\gamma c} \right)} \right) (\pi r_H^2) = \Phi_0 N = \frac{h}{2e} (3 \text{lobes} \times 2 \text{Positron Motion}) \quad (2)$$

$B = \mu_0 i / 2r_H$ is the minimum B inside the loop, and given r_H cancels out in eq.2, can be taken as a variational principle optimization of the energy B^2 .

Each of the 2 positron flux contributions around the circle ($N=2$). But each positron moves through all 3 $2P_{3/2} = N=2$ lobes. So doing the cancelations in eq.2:

$$\gamma(\mu_0/4)ec = (h/2e)(2 \text{positrons} \times 3 \text{lobes}). \quad (3)$$

So

$\gamma(\mu_0/4)ec = (h/2e)6$, But there already is a populated state (Hund's rule) $1S_{1/2}(\mu) = .1125 = \mu/P$ so we add it in (For example recall in the hydrogen atom that the 1S states fill before the 2P states.).

So:

$$\gamma = \frac{h}{2e} 6(1 + \mu) \frac{1}{\frac{\mu_0}{4} ec} \quad (\text{Note that 4 cancels the 4 in } \mu_0 = 4\pi \times 10^{-7} \text{ Wb-m/Amps.})$$

$$\gamma = \frac{2.0678 \times 10^{-15} (6)(1 + \mu)}{\pi \times 10^{-7} 1.6 \times 10^{-19} 3 \times 10^8} = \frac{1.2407 \times 10^{-14} \times (1.11255)}{1.5086 \times 10^{-17}} = \frac{1.38034 \times 10^{-14}}{1.5086 \times 10^{-17}} = 915 \quad (4)$$

We must add in the $3 \times .511 = 1.533$ for the 3 electrons

$$915 + 1.533 = 916.533$$

$2P_{3/2}$ at $r=r_H$ implies also twice our 2 positron γ result will be the proton mass.

$$2(916.533)m_e c^2 = 1.50087 \times 10^{-10} \text{ J} = 937 \text{ MeV}$$

Finally we must add that 1MeV binding energy between that μ and the (Fitzgerald contracted) net $+e$ positrons and electron (Fitzgerald contracted to a point Coulomb source) from axial frame of reference (sect.10.5) and get 938.23MeV.

Actual proton mass = 938.272MeV = m_p .

An exact answer!

Small C limit Finally Realized

Note we then we now have that small C limit from stable composite $3e$ because the positrons gain mass (**explaining the proton mass**) by their rapid motion contracting E field lines seen at the center area thereby (**explaining the strong force**).

m_p in equation 9.6 (and the rest of ch.8 is this 938 proton mass normalized to 1.

Also from appendix A7 scissors $r=r/\cos\theta$), eigenfunction perturbation (caused by object C in A7, gives the Fermi G).

In Ch.9 we perturb the Newpde $2P_{3/2}$ at $r=r_H$ state using a Frobenius series formulation. We then note that the Meisner effect $J=0$, $N=0$ Frobenius zero point energy 9.23 must add an extra e^+ to that μ^- to get the π^- case1 $J=0$ (Thus this later Frobenius solution ($J=0$, $N=0$ solution) formulation requires an (implicitly assumed) additional charge e^+ .)

Ortho State Eq.9.22 Zero Point Energy ϵ Implies Meisner Effect Nonzero Ortho States

$$m=1,0,-1. \gamma_\epsilon = \epsilon / \Delta\epsilon$$

The magnetic field in one of these protons is about 10^{11} T ($=\mu_0 i / (2r_H)$), so large that any spatially oscillating charge is going to be forced to induce a counter current that tries to cancel the change in flux produced by the charge motion (Faraday's law) relative to the proton. The Frobenius series method applied to the new pde has this $J=0$, $N=0$ zero point energy solution eq.9.22 SP hybrid state of the proton whose oscillation provides a Cooper pair oscillation counter current in that huge 10^{11} T field that cancels it out. So at close range there are many pions $\epsilon / \Delta\epsilon$. (up to 7) and more distant, where the B field drops you only need one: Hence we have just derived the multi pion interactions observed near the proton and resulting Yukawa force.

The ortho state with B orthogonal to A would not exist without this zero point SP state motion since the (SP hybrid so) induced P state spinor has a horizontal component so has a dot product with horizontal S_2 nonzero spinor (for ortho).

These two B fields (B_{\perp} and B_{\parallel}) are put into Paschen Back (eg., $m_s c^2 = u_B \gamma_e B(1+0+0+0)$)




Fig1

2 body positron-positron dynamics so ortho(s.c.b) and para(t) states on the $r=r_H$ shell near $\theta=90^\circ$. Paschen Back dynamics

Summary Of Para and Ortho States: two γ s: top view Para $\gamma=917$, Ortho side view $\gamma=\epsilon/\Delta\epsilon$
 S_{\perp} SP hybrid zero point energy eq.9.22: P state so $S_1 \bullet S_2 \neq 0$, ortho: $\gamma_{\epsilon m=\pm 1,0} B = (\epsilon/\Delta\epsilon) B = B_{\perp}$.
 $= (.06/.00058) 2.02 \times 10^{11} = 2.27 \times 10^{13} T = B_{\perp}$ (eg., $m_s = u_B B_{\perp} (1+0+0+0)$)

S_{\parallel} Flux quantization γ top S_{\parallel} view both positrons so $S_1 \bullet S_2 \neq 0$, para: $\gamma_{L=0} = \gamma B = B_{\parallel}$.
 $= 917(4) 2.02 \times 10^{11} = 7.5 \times 10^{14} T = B_{\parallel}$ (eg., $m_t = u_B B_{\parallel} (1+1+1+1)$)

7.6 The two rapidly moving positrons: $2 \otimes 2 = 3 \oplus 1$ List of 1 Para SB_{\parallel} and 3 Ortho SB_{\perp} States
 Here Thomas LS LST $\equiv -(L_{BL} * (S_{AL} \text{ or } S_{CL})) K \pm g_s$ perturbation is subtracted off the Paschen Back energy for both the SB_{\perp} and SB_{\parallel} cases.

SB_{\parallel} = State t non LS coupling para singlet state (the 1 single state in the $3 \oplus 1$ decomposition)
 $B_{\parallel} = 7.448 \times 10^{15} T, = B = \gamma (2.02 \times 10^{11})$

0° r_H
 $B_{\parallel} u_B (mLA + mSA + mLC + mSC) = PE$ LST PE-LST name Pauli Principle. L_{EM} S
 $L=0$ 1 + 1 + 1 + 1 173 0 173 t even stable Singlet **para**

SB_{\perp} State B total **triplet** b,c,s ground state u/d LS coupling triplet **ortho** state. LS coupling
 $B_{\perp} \approx 4.043 \times 10^{12} T$ $(\epsilon/\Delta\epsilon) 2.02 \times 10^{11} = 2.27 \times 10^{13} T$

90° $S_B = \pm 1$ r_{snf} $2P_{1/2} =$ at $r=r_H$. Here single $PE = \frac{1}{2} PE = \frac{1}{2} D$ bond $= \frac{1}{2}$; $D=2$. See sect.10.7

$B_{\perp} u_B (mLA + mSA + mLC + mSC) = PE$ LST \approx LST-PE name Pauli Principle
 $m=1$ 1 + 1 + 1 + 1 5790 1.5+1 (2) Ξ_b **ortho**

m=0	1 + 1 + 0 + 0	2471	1.5-0 (2)	Ξ_s	ortho
m=-1	1 + 1 + 0 + 0	1314	1.5-1 (2)	Ξ_s	ortho
Ground State $P_{u/d}=(1)938$		1	$P_{u/d}$	$2P_{3/2}$ & $2P_{1/2}$	

So a total of 4 states for two positrons (3ortho, 1para). 6 $2P_{3/2}$ states if you include the central electron. Since the proton is the core object for these states we can use the Frobenius solution Ch.9 perturbations below for these $r>r_H$ deviations from the spin 1 flux quantization $2\phi_E=2h/2e$ above sect.10.13 and Ξ . We get four multiplets of the three Ξ one P. Get ud. (Chapter 8). The above are also boson energy transitions analogous to the principle quantum number photon transition emissions of the hydrogen atom.

Other Ortho Consequences

We can reverse engineer this process by modeling a large decrease in the resulting strong magnetic field:

Neutron $2P_{1/2}$ $1-r_H/r_H$ for charge 0 (case 2 Ch.8) is homeomorphically mapped into $1-\varepsilon$ with added outside particle KE Meisner effect additional outside charge (reducing that r_H/r_H charge so preserving angular momentum a (and so KMQ) in the Kerr metric term $(a/r)^2$. Note the negative sign still indicates inside multibody charge is still 0.

Proton $2P_{3/2}$ $1+r_H/r_H$ (case 1 in Ch.8) is then homeomorphically mapped into $1+\varepsilon$ with added particle KE. The positive sign indicates nonzero internal multibody charge. See eq.B4.

For $2P_{3/2}$ $\kappa_{00}=(1-2\varepsilon)-\Delta\varepsilon-[(C_M/m_e)r]$. The starting point of PartII. (B3)

vector ω_p . So ultrarelativistic Thomas precession $=n3m_p = LS$ energy ω is subtracted off from Paschen Back energy ω_p . It adds to 0 in the ground state. Iso $L_{EM}=Nr_H\hbar\omega/c=rX[(EXH)/c](\pi r_H^2 T)$ angular momentum also cancels some of the total angular momentum of objects A,C and B.

For each ortho state we apply the Frobenius solution perturbation(Ch.9) . The next ortho value is $m_p=2$ (for s and later $m_p=4$ for c, $m_p=6$ for the b state) for the next ortho state.

Calculation Of ξ_1

We use the equation 1.2.7 energy normalization ($m_e=1$) for two reduced mass $2P_{3/2}$

ultrarelativistic positrons at $r=r_H$ with ansatz $\xi \rightarrow x_2$, in $\xi_0 \rightarrow 1$ in $\xi_1=\xi_3+\xi_2+\xi_0$. So $E=C_M\xi_1^2/\xi\sqrt{2} \rightarrow \frac{1}{2}\xi_1^2/\xi_1 = \frac{1}{2} / (x_2 + (2+\Delta)x + (1+\Delta)) =$ (partial fractions) $= \frac{1}{2} ((1(-1/\Delta))/(x+1)) + ((1(1/\Delta))/(x+1+\Delta)) =$ positron1 +positron2. So for $x \rightarrow 0$ then $\Delta=1/3684$ from the boosted magnetic flux calculation $2\gamma=3684$.

Composite 3e Paschen Back




Fig.8




Fig.2

Chapter 8 Frobenius Series Perturbation To Each Paschen Back State Introduction

Here we start with the ground state magnetic flux energy(u/d) set $m_b=1$, move on to the three orthos (s,c,b) with larger m_b s (Ξ) and finally to the very high para (t). We are actually perturbing the motions at r_H by these r in equation 9.5 and so are taking into account the constituents of the proton in this way.

Also there are then 6 magnetic flux quantization $2P_{3/2}$ states. Each flux quantization level has its own m_p and associated Frobenius solution. So we have ground state $m_p=1$, (appendixC 938Mev result) and excited states: $m_p=1.5=\Xi_s$, and also Ξ_c , Ξ_b , each having it's own Frobenius solution sets.

8.1 Solution to eq.2 Using Separability: Gyromagnetic Ratios And Low Energy Particles (energy<3GeV) Derived For ground state Magnetic Flux

$r \approx r_H$ Application: Gyromagnetic Ratios

After separation of variables the “r” component of equation 9 can be rewritten as:

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] F - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{J+3/2}{r} \right) f = 0 \quad (8.1)$$

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] f + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{j-1/2}{r} \right) F = 0. \quad (8.2)$$

Because the $\kappa_{00}=1-r_H/r$ is point source the object B ambient metric is local and so the vacuum is not infinite density (see also sect 6.11) as in the QED ambient metric which is homogenous.

Comparing the flat space-time Dirac equation to equations 8.1 and 8.2:

$$(dt/ds)\sqrt{\kappa_{00}}=(1/\kappa_{00})\sqrt{\kappa_{00}}=(1/\sqrt{\kappa_{00}})=\text{Energy}=E \quad (8.2a)$$

Using the above Dirac equation it is easiest to find the gyromagnetic ratios g_y for the spin polarized $F=0$ case. Recall the usual calculation of rate of the change of spin S gives $dS/dt \propto m \propto g_y J$ from the Heisenberg equations of motion. We note that $1/\sqrt{g_{rr}}$ rescales dr in

$\left(\sqrt{\kappa_{rr}} \frac{d}{dt} + \frac{J+3/2}{r} \right) f$ in equation 8.1. Thus to have the same rescaling of r in the second term we must multiply the second term denominator (i.e., r) and numerator (i.e., $J+3/2$) each by $1/\sqrt{g_{rr}}$ and set the numerator equal to $3/2+J(g_y)$, where g_y is now the gyromagnetic ratio. This makes our

equation 8.1 compatible with the standard Dirac equation allowing us to substitute the g_y into the standard $dS/dt \propto m \propto g_y J$ to find the correction to dS/dt .

Thus again:

$$\begin{aligned} [1/\sqrt{g_{rr}}](3/2 + J) &= 3/2 + Jg_y, \text{ Therefore for } J = 1/2 \text{ we have:} \\ [1/\sqrt{g_{rr}}](3/2 + 1/2) &= 3/2 + 1/2 g_y = 3/2 + 1/2 (1 + \Delta g_y) \end{aligned} \quad (8.3)$$

Then we solve equation 8.3 for g_y and substitute it into the above dS/dt equation.

S States: Recall ε and $\Delta\varepsilon$ and S states from eq. 6.4.13. These are zero point energy states (eq.9.22) that must also be the source of the Meisner effect canceling of those large B fields. Noting in equation 6.4.13 we get the gyromagnetic ratio of the electron with $g_{rr} = 1/(1 + \Delta\varepsilon/(1 + \varepsilon))$ and $\varepsilon = 0$ for electron. Thus solve equation 8.3 for $\sqrt{g_{rr}} = \sqrt{1 + \Delta\varepsilon/(1 + \varepsilon)} = \sqrt{1 + \Delta\varepsilon/(1 + 0)} = \sqrt{1 + 0.0005799/1}$. Thus from equation 8.3

$[1/\sqrt{1 + 0.0005799}](3/2 + 1/2) = 3/2 + 1/2(1 + \Delta g_y)$. Solving for Δg_y gives anomalous **gyromagnetic ratio correction of the electron** $\Delta g_y = .00116$

Going to higher energies (so $\varepsilon \neq 0$ in equation 8.3) we get the anomalous **gyromagnetic ratio correction of the muon**. From the momentum representation of eq.8.1,8.2:

$2P_{3/2}$ states: Recall the $2P_{3/2}$ states from chapter 3. Note also that k can be positive or negative since $4\pi k = Z_{00}$ in our Lagrangian with a positive k meaning at least one charge is not canceled. Therefore $1/g_{rr} = 1 \pm k/r + \varepsilon$ (using our Frobenius solution expansion near $r \approx r_H$ of eq.9.5 below multiply through by zero point energy, Meisner effect $(1 + \varepsilon/4)((1 + \varepsilon + \dots) \approx 1 + .08 = 1 + \varepsilon'$ so a pion mass is then added to the protons) from the \pm nature of Z_{00} . Therefore we have two cases from equation B3 at the boundary $r = k$

CASE 1	$1/g_{rr} = 1 + k/k + \varepsilon$	charge 1	(core case)
CASE 2	$1/g_{rr} = 1 - k/k + \varepsilon$	charge 0	(use m from case 1)

Note: ε (9.22) is required because it is the zpe here (like $\hbar\omega/2$ is the zpe of 1D SHM) external to the $3e$ region. So through the Fzero point energy araday's law Meisner effect pops up to cancel that huge $10^{14}T$ internal B field, hence the origin of the mesonic field. So the ε in case 1 and case II is the artifact of that large internal B field of section 8.1.

Also the effect of a zero charge is to make metric component $g_{00} (= 1/g_{rr})$ contribution zero in case 2. Thus the effect of *nonzero* charge is to increase the dimensionality by adding a metric component in eq.2. This provides the reason that Kaluza Klein theory (adding a 5th dimension) is so successful at injecting E&M into general relativity. But Kaluza Klein theory is not required here because finite C_M in eq.1.11 is really responsible for charge and E&M. 2D is sufficient as we showed in Chapter 1, eq.1.5. The extra 2D degree of freedom is associated with that extra real term $\delta\delta z$ in the amazing equation 1.6.

CASE 1: Plus $+k$, therefore is the proton + charge component. $1/g_{rr} = 1 + k/k + \varepsilon = 2 + \varepsilon$. Thus from equation 8.1, 8.2 $\sqrt{2 + \varepsilon}(1.5 + .5) = 1.5 + .5(g_y)$, $g_y = 2.8$ (8.4)

The gyromagnetic ratio of the proton (therefore that above $r \approx k$ stability was indeed proton stability as we concluded) $\text{mass} = m_p$. $dt/ds \sqrt{g_{00}} = 1/\sqrt{g_{00}} = E = m_p$

CASE 2: negative k , thus charge cancels, zero charge:

$$1/g_{rr} = 1 - k/k + \varepsilon = \varepsilon \quad \text{Therefore from equation 8.3 and case 1 } 1/g_{rr} = 1 + k/k + \varepsilon$$

$$\sqrt{\varepsilon} (1.5 + .5) = 1.5 + .5(gy), \quad gy = -1.9, \quad (8.5)$$

the **gyromagnetic ratio of the neutron** with the other charged and neutral hyperon magnetic moments scaled using their masses by these values respectively.

Chapter 9

The composite 3e Energies For particle energy <3GeV Derived Using Frobenius Series Solution (at first Paschen Back m_p level) Perturbation)

9.1 Series Solutions ψ Ansatz Near $r \approx r_H$

$m_p = 1$ here. m_p determined from Paschen Back energy level (next Paschen Back level $m_p = 2$ for s)

Recall equations 8.1, 8.2:

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] F - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{j+3/2}{r} \right) f = 0$$

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] f + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{j-1/2}{r} \right) F = 0$$

Recall from the previous section $g_{00} = 1 - k/r - (\varepsilon + \Delta\varepsilon)$. Also **recall our Dirac doublet** (equation 7) must have a left handed zero mass component will be called case 1 and case 3 respectively below. Also we need the equivalent of the singlet equation 2 is our below case 2. Also in equation 2 at $r = r_H$ the eigenvalue is $\Delta\varepsilon + \varepsilon + 1 = 2m_p$ for that principle quantum which then must be the same for the $2P_{3/2}$ state. Here we write out the left handed Dirac Doublet Eq.2 in the general representation of the Dirac matrices. Also recall from chapter 8 that the $2P_{3/2}$ state (and its sp^2 hybrid) for this new electron Dirac equation gives a azimuthal trifolium, 3 lobe shape and thus a $\lambda/3$ spherical harmonic wavelength so that for covalent bonding $r' \approx r_H/3$ in $\kappa_{00} = 1 - r'/r$. This $\lambda/3$ also is used to calculate P wave scattering (called “jets” by quark people.)

To use the f & F components of the equation 8.1, 8.2 Dirac equation we write the Dirac equation for free particle motion along the symmetry axis z (r=ratio of momentum to energy) to find the chirality of the components in the general representation of section 1.6. We then compare this z motion free particle Dirac equation eigenfunction structure with radial component structure to arrive at a sense of which components of the radial equation are left handed and which aren't. This step is a little more complicated here because we are not using the chiral representation of the Dirac matrices, but the standard representation instead. In any case given that the electron is positive energy, then (as we see below) for the positron -E gives left handed f and F implying that this object *must* have a positive charge since this left handedness(doublet, Ch.3) results from the fractalness (There is a corresponding argument for G and g). The proton indeed is positive charged. So:

$$\left[-\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] g - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{J+3/2}{r} \right) G = 0. \rightarrow \mu c^2 u_1 + c p u_3 - E p u_1$$

$$\left[-\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] G + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{J-1/2}{r} \right) g = 0 \rightarrow c p u_1 - \mu c^2 u_3 - E p u_3$$

$$\left[-\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] f - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{J-1/2}{r} \right) F = 0 \rightarrow \mu c^2 u_2 - c p u_4 - E p u_2$$

$$\left[-\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] F + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{J+3/2}{r} \right) f = 0 \rightarrow -c p u_2 - \mu c^2 u_4 - E p u_4$$

where to get correspondence from these two Dirac equation structures

$$\text{we see that at } +E: u^R = \begin{pmatrix} 1 \\ 0 \\ r \\ 0 \end{pmatrix} = g, \quad u^L = \begin{pmatrix} 0 \\ 1 \\ 0 \\ -r \end{pmatrix} = f; \quad -E: \text{No } (v^R = \begin{pmatrix} r \\ 0 \\ 1 \\ 0 \end{pmatrix} \text{ here}), \quad v^L = \begin{pmatrix} 0 \\ r \\ 0 \\ 1 \end{pmatrix}$$

$$= F, \text{ Note in general (with } r \approx 0) \text{ here: } \begin{pmatrix} ig \\ if \\ G \\ F \end{pmatrix} = \begin{pmatrix} u^R \\ u^L \\ v^R \\ v^L \end{pmatrix} = \Psi. \text{ So we have the}$$

solution that in the standard representation of the left handed doublet is given by F and f only for -E of the electron (here a positron needed below for + proton hadron excited states) at the horizon. Dirac matrices

$$\text{So for the left handed doublet: } \begin{pmatrix} F \\ f \end{pmatrix}_L \text{ we have respectively } \begin{matrix} q \pm, m \neq 0, -E, \text{ for } F \\ q = 0, m = 0, +E, \text{ for } f \end{matrix} \quad (9.4)$$

Or more succinctly equation 2 in the Dirac doublet form implies in section B2: Note our postulate implies $C \rightarrow 0$ so we are on the dr axis thus $dt' = 0$ so $dt'^2 = (1 - r_H/r) dt^2$ (sect.0.1 of Ch.1). Thus $r = r_H = k$ is a stable point since the clock stops since $dt' = 0$ and this is the Meisner effect formalism for canceling out that huge B field at a distance and also making it so the protons mass is mp, and not much larger. .

$$\text{CASE 1} \quad 1/\kappa_{rr} = 1 + k/k + \epsilon = 1 + r_{HM+1}/r + r_{HM}/r + \epsilon \quad (\text{core case})$$

$$\text{CASE 2} \quad 1/\kappa_{rr} = 1 - k/k + \epsilon = 1 + r_{HM+1}/r + r_{HM}/r + \epsilon$$

Normalize out $1 + r_{HM+1}/r$. That just divides by 2 since we (at r) are already near the event horizon

$$\text{CASE 1} \quad 1/\kappa_{rr} = 1 + k/k + \epsilon = 1 + r_{HM}/r + \epsilon \quad \text{charge 1} \quad (\text{core case})$$

$$\text{CASE 2} \quad 1/\kappa_{rr} = 1 - k/k + \epsilon = 1 + r_{HM}/r + \epsilon \quad \text{charge 0}$$

So if $|r_{HM+1}/r| = |r_{HM}/r|$ (use m from case 1) then negative r_{HM}/r means zero charge (so $r_{HM+1}/r = r_{HM}/r$ so charge sources cancel out) and positive means charged. (see also above sect.B2).

Note in sect.1.5 we can have a zero and nonzero charge in the 3rd quadrant (where $dt = dr$) massive Proca boson case given the possibilities in sign we have for $\pm \epsilon'/2$ in $((-\epsilon/2) \pm \epsilon'/2) dr - ((-\epsilon/2) \pm \epsilon'/2) dt$.

In the first quadrant $ds = 0$ again (section 1.4) so they have to add to zero. $+dr + \epsilon/2 + dt - \epsilon/2$ and $-dr - \epsilon/2 - dt + \epsilon/2$ solutions. Multiply the second equation by -1, then add the two resulting equations, then divide by 2 and get $dr + \epsilon/4 \pm \epsilon/4 + dt - \epsilon/4 \pm \epsilon/4$ so that $\epsilon/2 \rightarrow \epsilon/2 \pm \epsilon/2$. So we multiply each of the two ds^2 cases (above $|dr + dt|$ discussion) by its own dz , each with its own $\kappa_{rr} = 1/(1 - \epsilon/r) \rightarrow 1/(1 - (\epsilon/2 \pm \epsilon/2)/r)$ (sect.4.7) implying 2 charges $\epsilon/2 - \epsilon/2 = 0$, $\epsilon/2 + \epsilon/2 = \epsilon$ and so two Proca equation massive W, Z .

See B2. .See above B2:

$$\text{CASE 1} \quad 1/g_{rr} = 1 + k/k + \epsilon \quad F \quad \text{charge 1, } m=1 \quad (\text{core case}) \quad 2P3/2$$

$$\text{CASE 2} \quad 1/g_{rr} = 1 - k/k + \epsilon \quad F \quad \text{charge 0, } m \text{ from case 1) } 2P1/2$$

$$\text{CASE 3} \quad \quad \quad f \quad \text{charge 0, } m=0$$

We solve these equations only near $r \approx r_H$ since that is where the stability is to be found (and also fortunately were these equations are *linear* differential equations). Thus our first step is to expand $\sqrt{g_{rr}}$ about this radius and drop the higher order terms.

The Frobenius series solution method can now be used to solve equations 8.1 and 8.2 at $r \approx r_H$. See for example Mathematical Methods of Physics, Arfken 3rd ed. Page 454. First we solve the f in equation 8.1, plug that into equation 8.2 and then have an equation in only F . There we substitute a series solution ansatz $F = \sum a_n r^n$ in the resulting combined equations. We can then separate out the results into coefficients of respective r^n and get recursion relations that will give us series that must be terminated at some N . Note the energy Eigenvalue 'E' will be in this series as $dt/ds \sqrt{g_{00}}$ so we can then solve for the mass energy of these hadrons at specific J . We will need an indicial equation for the first term to start out this process. Also in this Frobenius solution method 'n' turns out to be a multiple of $1/2$ and the series must start at $n = -1$. Finally to get the charge zero case the charged case must be done first and its constant masses used in the uncharged state calculations.

Here in Ch.9 we perturb the Newpde $2P_{3/2}$ at $r=r_H$ state using a Frobenius series formulation. We then note that the Meisner effect $J=0$, $N=0$ Frobenius zero point energy 9.23 must add an extra e^+ to that μ^- to get the π^- case1 $J=0$ (Thus this later Frobenius solution ($J=0$, $N=0$ solution) formulation requires an (implicitly assumed) additional charge e^+ .)

9.2 CASE 1 charged: Excited States for F, $m \neq 0$, $q \pm 2P_{3/2}$

Again case 1 is one of the equation 8.1 possibilities. Therefore let $R=k_H-r$, $r < R$ (for stability) we can write in 8.1:

$$\sqrt{K_{rr}} = \frac{1}{\sqrt{1+\frac{r_H}{R}+\varepsilon}} \approx \frac{\sqrt{R}}{\sqrt{R+r_H+R\varepsilon}} = \quad (9.1)$$

$$\frac{\sqrt{r_H-r}}{\sqrt{r_H-r+r_H+(r_H-r)\varepsilon}} = \quad (9.2)$$

$$\frac{\sqrt{r_H-r}}{\sqrt{r_H(2+\varepsilon)-r(1+\varepsilon)}} = \quad (9.3)$$

$$\left(\frac{1-\varepsilon}{4}\right) \frac{\left[1-\frac{r}{2r_H}+\frac{r^2}{8r_H^2}+\dots\right]}{\sqrt{2}} = \quad (9.4)$$

$$\left(\frac{1-\varepsilon}{\sqrt{2}}\right) \left(1 - \frac{r}{4r_H} + \frac{3r^2}{32r_H^2} - \dots\right) \approx \frac{1-\frac{r}{4r_H}}{\sqrt{2}} \quad (9.5)$$

Note **taking the first term of this Taylor expansion of the square root makes this an approximation (<2GeV.)**. Note that including the above $1 \pm \varepsilon/4$ the compensating $(1 \pm \varepsilon/4)$ in the next r term has the effect of a multiplying the derivative terms by $1 \pm \varepsilon/4$. This rescales r to allow us to still say that the stable boundary is still at r_H . Thus we could use it to also rescale t in the first term of equations 8.1 and 8.2 or note that $(1 \pm \varepsilon/4)(1 \pm \varepsilon) = 1 \pm 5/4\varepsilon$ thus renormalizing $1 \pm \varepsilon$ to $1 \pm 5/4\varepsilon = 1 \pm \varepsilon'$ everywhere. Also the $3r^2/32r_H^2$ terms must be included. We drop these perturbative terms until the end. Therefore substituting in equation 9.5 we find that equation 8.1 reads:

$$\sqrt{K_{rr}} = \frac{1}{\sqrt{1+\frac{r_H}{R}+\varepsilon}} \approx \frac{\sqrt{R}}{\sqrt{R+r_H+R\varepsilon}} = \quad (9.1)$$

$$\frac{\sqrt{r_H-r}}{\sqrt{r_H-r+r_H+(r_H-r)\varepsilon}} = \quad (9.2)$$

$$\frac{\sqrt{r_H-r}}{\sqrt{r_H(2+\varepsilon)-r(1+\varepsilon)}} = \quad (9.3)$$

$$\frac{\left(1-\frac{\epsilon}{4}\right)}{\sqrt{2}} \left[\frac{1-\frac{r}{2r_H}+\frac{r^2}{8r_H^2}+..}{1-\frac{r}{4r_H}+\frac{r^2}{16r_H^2}+..} \right] \quad (9.4)$$

$$\left(\frac{1-\frac{\epsilon}{4}}{\sqrt{2}}\right) \left(1 - \frac{r}{4r_H} + \frac{3r^2}{32r_H^2} - ..\right) \approx \frac{1-\frac{r}{4r_H}}{\sqrt{2}} \quad (9.5)$$

Therefore

$$f = -\hbar c \frac{\hbar \epsilon}{(E-m_p)} \left[\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} - \frac{\left(1+\frac{r}{r_H}\right)\left(j-\frac{1}{2}\right)}{r_H} \right] F \quad \text{substituting into}$$

therefore

$$[E + m_p]F - \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{J+\frac{3}{2}}{r_H-r} \right) f = 0 \quad (9.7)$$

We find solving for f and substituting back in:

$$\begin{aligned} [E + m_p]F - \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{\left(1 + \frac{r}{r_H}\right)(J + 1.5)}{r_H} \right) * \\ \frac{\hbar c}{E - m_p} \left(-\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{\left(1 + \frac{r}{r_H}\right)\left(J - \frac{1}{2}\right)}{r_H} \right) F = (E + m_p)F + \\ \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(-\left(1 - \frac{r}{4r_H}\right) \frac{d}{4r_H\sqrt{2}dr} + \left(1 - \frac{r}{4r_H}\right)^2 \frac{d^2}{\sqrt{2}r^2} - \frac{\left(1 - \frac{r}{4r_H}\right)\left(j - \frac{1}{2}\right)}{r_H^2} \right) F \\ + \frac{(\hbar c)^2}{E - m_p} \left(\left(1 + \frac{3r}{4r_H}\right)(j + 1.5) \frac{d}{\sqrt{2}r_H dr} - \frac{\left(1 + \frac{r}{r_H}\right)^2(j + 1.5)\left(j - \frac{1}{2}\right)}{r_H^2} \right) F = \\ \left([E + m_p] + \left[\frac{(\hbar c)^2}{E - m_p} \left(-\frac{\left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^2} - \frac{\left(j - \frac{1}{2}\right)}{\sqrt{2}r_H^2} \right) \right] \right) F + \\ \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(-2\sqrt{2} \frac{\left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^3} + \frac{\left(j - \frac{1}{2}\right)}{4r_H^3} \right) rF + \\ \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\left(\frac{-1}{4r_H\sqrt{2}} \right) + \left(j + \frac{3}{2} \right) \frac{1}{r_H} \right) \frac{dF}{dr} + \\ \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{1}{r_H^2 16\sqrt{2}} + \left(j + \frac{3}{2} \right) \frac{3}{4r_H^2} \right) r \frac{dF}{dr} \\ \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{1}{\sqrt{2}} \right) \frac{d^2 F}{dr^2} + \end{aligned}$$

$$\frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{-1}{2\sqrt{2}r_H} \right) r \frac{d^2 F}{dr^2}$$

Here $r=2r_H$ is a regular singular point. Next substitute in $F = \sum_n a_n r^n$ with again half integer n allowed as well:

$$\sum_M^N \left([E + m_p] + \left[\frac{(\hbar c)^2}{E - m_p} \left(-\frac{(j+\frac{3}{2})(j-\frac{1}{2})}{r_H^2} - \frac{(j-\frac{1}{2})}{\sqrt{2}r_H^2} \right) \right] \right) a_n r^n + . \quad (9.8)$$

$$\sum_M^N \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(-\frac{2\sqrt{2}(j+\frac{3}{2})(j-\frac{1}{2})}{r_H^3} + \frac{(j-\frac{1}{2})}{4r_H^3} \right) a_{n-1} r^n + . \quad (9.9)$$

$$\sum_M^N \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(-\frac{1}{4\sqrt{2}r_H} + \frac{(j+\frac{3}{2})}{r_H} \right) (n+1) a_{n+1} r^n + . \quad (9.10)$$

$$\sum_M^N \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{1}{16\sqrt{2}r_H^2} + \frac{3(j+\frac{3}{2})}{4r_H^2} \right) n a_n r^n + \quad (9.11)$$

$$\sum_M^N \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{1}{\sqrt{2}} \right) (n+2)(n+1) a_{n+2} r^n + . \quad (9.12)$$

$$\sum_M^N \frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{-1}{2\sqrt{2}r_H} \right) (n+1) n a_{n+1} r^n = 0. \quad (9.13)$$

Note from equation 9.12 that this series diverges. To terminate the series we now take 9.8 and 9.11 together and 9.10 and 9.13 together (since they have the same a_n). For example combining the equation 9.8 and 9.11 terms

$$\left((E + m_p) + \left[\frac{(\hbar c)^2}{E - m_p} \left(-\frac{(j+\frac{3}{2})(j-\frac{1}{2})}{r_H^2} - \frac{(j-\frac{1}{2})}{\sqrt{2}r_H^2} \right) \right] \right) +$$

,

$$\frac{(\hbar c)^2}{(E - m_p)\sqrt{2}} \left(\frac{1}{16\sqrt{2}r_H^2} + \frac{3(j+\frac{3}{2})}{4r_H^2} \right) n.$$

Replacing the normalization $m_p \rightarrow m_p(1 \pm \varepsilon)$ (from section 4.8):

$$(E^2 - m_p^2) + \left(-\frac{(j+\frac{3}{2})(j-\frac{1}{2})}{r_H^2} - \frac{(j-\frac{1}{2})}{\sqrt{2}r_H^2} \right) - \frac{1}{\sqrt{2}} \left(\frac{1}{16\sqrt{2}r_H^2} + \frac{3(j+\frac{3}{2})}{4r_H^2} \right) N = 0$$

Therefore after rearranging:

$$E = \sqrt{m_p^2 + \frac{1}{r_H^2}(j^2 + 1.7071j - 1.10355 - (J.5303 + .8269)N)} \quad (9.14)$$

We have for a general Laurent series ansatz:

$$.. + a_{-1}r^{-1} + a_{-1/2}r^{-1/2} + a_0r^0 + a_{1/2}r^{1/2} + a_1r^1 + .. = F$$

Note also that equations 9.8-9.13 imply that the coefficients of a given r^n are independent. Thus adding together the coefficients of r^n for equations 9.8–9.13 at a given n :

$$9.9(j^{-1/2})a_{n-1} + (9.8+9.11)a_n + (9.10+9.13)(n+1)a_{n+1} + 9.12(n+2)(n+1)a_{n+2} = 0 \quad (9.15)$$

Method of Solving Equation 9.15

For the outside observer an $F=0$ finite boundary condition at infinity applies for flat vacuum value $n=0$, $j=1/2$ and for r^0 , $r^{-1/2}$, r^{-1} and for complete vacuum for $N=0$, $J=0$.

Here then the generalized Laurent series $.. + a_{-1}r^{-1} + a_{-1/2}r^{-1/2} + a_0r^0 + a_{1/2}r^{1/2} + a_1r^1 + .. = F$ reduces to $.. + a_{-1}r^{-1} + a_{-1/2}r^{-1/2} + a_0r^0 = F$. Thus either set $9.9(j^{-1/2})a_{n-1} = 0$ or $(9.10+9.13)(n+1)a_{n-1} + 9.12(n+1)(n+1)a_{n+2} = 0$ separately in eq.9.15 or set both equal to zero:

$J=1/2$, sets eq.9.9=0

- 1) $N=-1$, in equation 9.14 gives mass eigenvalue for Ξ
Exact solution for all possible a_n , sets none of them to zero.
- 2) $N=0$, in equation 9.14 gives mass eigenvalue for *nucleon*. $dr^0/dr=0$ so all derivative of F terms are then zero and this solution applies inside as well.
 $N=0$ flat $J=0$ allowed flat vacuum gives π^\pm and with free e , $j=1/2$ muon.
- 3) $N=-1/2$, in equation 9.14 gives mass eigenvalue of two Σ s since a plus and minus square root of r .

These $9.9=0$ cases have case 2 zero charge representations as well.

$N=-1$, Principle QM number Also $a_{-2}=0$

- 1) $J=0$, in equation 9.14 gives mass eigenvalue for K
- 2) $J=1$, gives deuteron mass eigenvalue (bonding) given $N=0, J=0$ fills first (i.e., pion). Thereafter use nuclear shell model-Schrodinger equation many body techniques with these nonrelativistic lobes with this (bound state) force acting like a outer layer surface tension, finite height square well potential. Get an aufbau principle that then gives the D,F,G,..nuclear shell model states. Alternatively can fill that first S state in with free $1S_{1/2}$ (next state to filled state) and we have $j=3/2$ Ω^- filling in some (i.e., uds) of the $2P_{3/2}$ states (see Ch.9) and thereby also deriving from first principles Gell Man's 1963 eight fold way for hyperon eigenvalue classification (to finish that effort need case II zero charge and case III Λ_0 as well). M_p is replaced by 2 in c hyperons, by 4 for b hyperons as indicated in fig. 16-1 for how to fill in the cbt 2P harmonic states given the requirement to use r^2 then.

Also, to include higher order r expansion term effects in equation 9.5 we must include those perturbative $1+\varepsilon/4$ and $3r^2/32k_H^2$ contributions which gives a $n(n-1)/6.4$ added to the “ n ” term component inside the radical of equation 9.14.

In our new pde $\delta J=0$ through LS spin-orbit coupling so the three spin $1/2$ s and the $L=1$ add to a minimum. $1-1/2-1/2+1/2=1/2=S$ for the proton with possible Pauli principle non $S=1/2$ possibilities for larger mass eigenvalue.

Details of Above Solutions for Case 1

Thus besides the ground state ($N=0$ $F_{\text{groundstate}} = \sum a_n r^n = a_0 r^{-0} = F_0$ proton) we have the two solutions:

$$F_{N=-1} = \sum a_n r^n = a_{-1} r^{-1} = F_1, j = 1/2, 0, \quad F_{N=-1/2} = \sum a_n r^n = a_{-1/2} r^{-1/2} = F_2. \text{ For } j = 1/2, 0.$$

Note the energy eigenvalues (E) can be found from the solution to equation 9.14 and $k_H=1$ with $E=1=938\text{MeV}$. Thus

$N=0, j=1/2$ then 9.14 gives +**Nucleon** (ground state) mass eigenvalue. Note that for the $N=0$, (with $J=1/2$ and also $J=0$ in section 9.5) ground state that the charge density is uniform (i.e., $\rho = K \propto r^0$) for $r < k$.

$N=-1/2, j=1/2$ **two** valued because of the *two* square root solutions. Equation 9.14 then gives Σ^\pm (charged sigma particle) **1184MeV particles, F_2 eigenfunction(s). Actual 1189MeV**

$N=-1, j=1/2$ gives *one* charged Ξ particle. Therefore the energy from equation 9.14 is **1327 MeV (actual 1321), F_1 eigenfunction.**

Case 2 and case 3 give the neutral hyperons and Λ_0 respectively (see case 3 below).

9.5 Nucleon Wavefunction: $J=1, q \neq 0, N=-1$ of Case 1

Here we recall case 1, section 9.3 above and compute energy eigenvalues for $J=0$ and $J=1$. again using equation 9.14 in case 1.

$J=0$

$N=-1, j=0$ **$E=490\text{ MeV}$ from equation 9.14 case 1. K^\pm .** Substitute into strangeness equation 9.34 case 1 we obtain strangeness =1.

$N=0, j=0$ then **from equation 9.14 $E=139.7\text{ MeV}$** (9.22)

case (note again $m=1+\varepsilon=1.061$ in 9.14 for outside). This is the nontrivial F zero point energy (and so has a fundamental SP hybrid state harmonic) for $r < k = \varepsilon$ at $r=r_H$. since the square root in equation 20.1 becomes imaginary then. Thus the mass of π^\pm is now the vacuum (e.g., note $F \propto r^0$ for $N=0$ here) ε' at $r \approx k$ explaining why this fundamental harmonic result for π is used in all the successful nuclear force theories such as in the Skyrminion Lagrangian for example. Note that:

$$m_{\pi^\pm} = 139\text{MeV} = 1.3(105.6\text{MeV}) = 1.3\varepsilon = .08\varepsilon'$$

$N=-1, J=1$ case 1. Recall for $J=1$ we have $\psi \propto r \sin\theta \propto Y^1_1(\theta, \phi)$ double lobe $\psi^* \psi$ along the z axis: From equation 9.14 we find with these inputs that **$E=1867\text{MeV}$** (9.23)

implying that (because $E \sim 2m_p$ and $J=1$) this eigenstate is responsible for the spin 1 deuteron (state). The $L=1, 2P$ state solution(s) are symmetric and so of the form $(1/\sqrt{2})(\psi_1 \psi_2 + \psi_2 \psi_1) = \psi_s$ and have positive parity even if the $2P$ ψ_1 and ψ_2 each has negative parity. The Deuteron thus has + parity (Enge, 1966).

Recall if we include the background metric in eq.6.4.11 $\kappa_{00} = 1 + r_H/r + 2\varepsilon' + \Delta\varepsilon$ and

$\kappa_{rr} = 1/(1 + r_H/r + \Delta\varepsilon)$. So rescaling $r \rightarrow r - \varepsilon' = r'$ for r near r_H allows us to use our above solutions

again. So in equation 8.1 $(1/\sqrt{\kappa_{rr}})\psi = 1/\sqrt{(1 + r_H/r' + \Delta\varepsilon)}\psi \approx 1/\sqrt{(1 + r_H/r)}\psi + (\varepsilon/2)\psi$. Note if we again

rescale our numerator $J=1 \rightarrow 1+(\varepsilon'/2)^2$ so that we have perturbed our Y_1 spherical harmonic with a $(\varepsilon'/2)Y_2$ giving a measure of the oblate, non spherical structure (e.g quadrupolar ψ_D and higher. $\varepsilon'/2 \approx .04$ from 9.22 therefore the nonspherical component of ψ is approximately 4% of the total ψ and is often called the tensor component of the Deuteron eigenstate (Enge, 1966). This simplest multiparticle state represents the *deuteron* state and this is then the explanation for the deuteron tensor component of the nuclear force.

Also the energy of the Deuteron is given just outside the r_H boundary (so $\varepsilon' \rightarrow i\varepsilon$ in 6.4.11) by $E_D = \text{Re} 1876/\sqrt{k_{00}} = \text{Re} 1876/\sqrt{(1+i\varepsilon') + \dots} = 1876(1-i\varepsilon'/2 + (3/8)(i\varepsilon')^2 + \dots)$. So the added real term due to the ε' is equal to $1876(3/8)\varepsilon'^2 = 1876(3/8)(.08)^2 = 4\text{MeV}$. In free space $\varepsilon'=0$ and just outside the nucleus it gives this contribution to the Deuteron energy. Thus this $(3/8)\varepsilon'^2$ is the binding energy of the Deuteron.

Note from the equation 9.15 discussion for $N \text{ not } -1$ we can only use $J=1/2$ and $J=3/2$ thus are restricted for two particles to S and P states (i.e. $1/2 + 1/2 = 1$) which then gives us the hyperons. For $N=-1$ we can use other J and can thereby construct large nuclei.

The multinucleon nuclei really are the solutions of the indicial equations of 9.15.

Recall in the shell model a hard shell nuclear outer wall is assumed with free space oscillations allowed inside this shell. The solutions to the Schrodinger equation are then spherical Bessel functions with corrections for spin orbit interaction, finite well height and tapered wells (Herald Enge, Introduction To Nuclear Physics, P.145). In any case an infinite mean free path for these oscillations is assumed to exist inside this shell. So how can there be an infinite mean free path inside this extremely high mass density region?

In that regard the above 2, $J=1$, $N=-1$ 2P deuteron state can also be viewed as yet another Bogoliubov pairing interaction (such as in the SC section 4.5) giving this infinite mean free path of the electron pairs comprising a pion acting as a Cooper pair, just as in SC In the context of the section 4.5 pairing interaction model $A(dv/dt)/v^2$ is no longer as small but dv/dt becomes very large to due to the ultrarelativistic motion of the electrons inside the nucleons. In any case this infinite mean free path for these oscillations (recall Cooper pairs have an infinite mean free path) is thereby explained here as a new type of superconductivity.

Spin Orbit Interaction In Shell Model

Recall the derivation of the shell model from first principles in section 6.12. If equal numbers of Neutrons and Protons gyromagnetic ratios then $g_P - g_N = 2.7 - 1.9 = .8$.

Since more neutrons in heavier elements: $(1/1.1)(.8) = .7$.

$R = r_H \equiv 1/2$ Fermi measured from singularity at $1 - 1/2 = 1/2$.

From $2P_{3/2}$ at $r=r_H$ Fitzgerald contraction discussion in section 2.2: $r \rightarrow R = 1/2(1 - 1/2) = 1/4$ Fermi \equiv

$R_V(r - r_H)$ so $R_V(r - r_H) \rightarrow Kr$. From Ch1, sect 4.16 $V = 1/(r - r_H)$. Spin orbit interaction =

$a_0^2(1/r)(\partial V/\partial r)(s \cdot L) =$

$$a_0^2 \frac{1}{R_V(r - r_H)} \frac{\partial V}{\partial (R_V(r - r_H))} (s \cdot L) = \frac{.7}{R_V(r - r_H)} \left(\frac{-1}{(R_V(r - r_H))^2} \right) (s \cdot L) =$$

$$= .7(4^3)(s \cdot L) \frac{1}{r} \frac{\partial V}{\partial r} = .7(64)(s \cdot L) \frac{1}{r} \frac{\partial V}{\partial r} = a_0^2 \frac{1}{r} \frac{\partial V}{\partial r} (s \cdot L) =$$

45*E&M spin orbit interaction.

Thus the $a_0=1$ Fermi. Thus the nuclear spin-orbit interaction is much larger than the E&M spin orbit interaction because the nucleons are much closer to r_H than to $r=0$ and the Fitzgerald contraction of the nucleon $2P_{3/2}$ state is on the order of $1/2$.

At close range there are higher energies available so the 4mev (=be) in equation 9.3 (if we include r^2 contributions) becomes the binding energy for the deuteron in $g_{00}=1-k/r+be$ in 8.1 particles, F_2 eigenfunction(s). Actual 1189Mev

$N=-1, j=1/2$ gives one charged Ξ particle. Therefore the energy from equation 9.14 is 1327 Mev (actual 1321), F_1 eigenfunction Ξs the fundamental structure for $m=1.5$. So we reapply the analysis all over again for $mp>1.5$ instead of 1.

Case 2 and case 3 give the neutral hyperons and Λ_0 respectively (see main Frobenius series solution paper).

The multinucleon nuclei are the solutions of the indicial equations of 9.15.

Recall in the shell model a hard shell nuclear outer wall is assumed with free space oscillations allowed inside this shell. The solutions to the Schrodinger equation are then spherical Bessel functions with corrections for spin orbit interaction, finite well height and tapered wells (Herald Enge, Introduction To Nuclear Physics, P.145). In any case an infinite mean free path for these oscillations is assumed to exist inside this shell. So how can there be an infinite mean free path inside this extremely high mass density region?

In that regard the above 2, $J=1, N=-1$ 2P deuteron state can also be viewed as yet another Bogoliubov pairing interaction (such as in the SC section 4.4) giving this infinite mean free path of the electron pairs comprising a pion acting as a Cooper pair, just as in SC In the context of the section 4.5 pairing interaction model $A(dv/dt)/v^2$ is no longer as small but dv/dt becomes very large to due to the ultrarelativistic motion of the electrons inside the nucleons. In any case this infinite mean free path for these oscillations (recall Cooper pairs have an infinite mean free path) is thereby explained here as a new type of superconductivity.

Particle Lifetimes

Recall from section 1.1: $\kappa_{00}=1-r_H/r$ so $r-r\kappa_{00}=r_H$ analogous to $dr-ct\kappa_{00}=ds$ so $r_H=ds\equiv|dZ|$. From section 6.7 there are three Dirac equation contributions with one being the ultrarelativistic m_v contribution. For that contribution we put Dirac αs into $dr+idt=dZ$ the free space Dirac equation. Dividing by ds gives mass on the right side in that Dirac equation. Because the motion of the $m_v=1eV$ (Ch.3) particle is ultrarelativistic in these hadrons we apply figure 1-1 $dr=dt$ so $\theta=45^\circ$ and so $dZ/ds=e^{i\pi/4}dr/ds$ for the ultrarelativistic m_v (on earth contribution of Ch.3). Note that $(e^{i\pi/4})^2=i$. We add another contribution (for spin $1/2, N=-1$) to get zero charge case II below. For added $2P_{1/2}$ ($K, \pm\pi$ mesons) there are 3e in r_H below (sect.10.3). Thus we obtain:

$$\text{hyperons, Kaons and } \pm\pi: \quad e^{i\pi/4}2e^2/m_v c^2 = e^{i\pi/4}r'_H = R_H$$

Recall that domain $r=r_H$ was the most stable, the proton state. This stability condition can be restated in terms of excess energy above the proton rest mass. Next substitute this m and ultrarelativistic m_v in the r_H in equation 9.14 with this r'_H in the relativistic solution of equation 2 described in Ch.1,sect.1.

$$E = \sqrt{m_p^2 + \frac{1}{R_H^2} \left(j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N \right)}$$

$$\approx m_p \left(1 + \frac{(e^{i\pi/4})^2 (j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N)}{2m_p^2 r_H'^2} \right).$$

Add to above to 9.14 result to get for the total energy:

$$m_p \left(1 + \left(\frac{e^{i\pi/4}}{r_H'} \right)^2 + \left(\frac{1}{r_H} \right)^2 \right) \frac{(j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N)}{2m_p^2}$$

Plug $(\hbar c/e^2)^2 = (1/\alpha)^2$ back in eq.8.1 and normalize $m_v c^2$ to 1/hz with 1/h. Next plug into the time propagator e^{iHt} and get for the r_H' (decay) term:

$$\begin{aligned} &= \exp i \left(\left((m_p c^2 / h) + (e^{i\pi/4})^2 (m_v c^2 / h) \frac{m_v}{m_p} (j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N) \right) \left(\frac{\hbar c}{2e^2} \right)^2 \right) t \\ &= \exp i \left(\left((m_p c^2 / h) + i (m_v c^2 / h) \frac{m_v}{m_p} \frac{i(j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N)}{(2\alpha)^2} \right) \right) t \quad (9.23) \\ &= \exp i \left((m_p c^2 / h) + i\Delta \right) t \text{ giving hyperon, Kaon, } \pm\pi \text{ decay times.} \end{aligned}$$

The second term Δ is also the excess mass above the proton mass.

For neutrons (939Mev) the excess mass above the proton mass (938Mev) is $m_p/1000$ and $R_H \rightarrow 1000R_H$, $\Delta \rightarrow \Delta'$

$$E^2 = m_p^2 + \frac{1}{1000^2} \frac{1}{(1000R_H)^2} (j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N)$$

gives the neutron decay time.

For m_μ muons $j=1/2$, $N=0$ and the excess mass is $m_p/8.87 \equiv m_\mu$.

$$E^2 = m_p^2 + \frac{1}{8.87^2} \frac{1}{(8.87R_H)^2} (j^2 + 1.7071j - 1.10355 - (j.5303 + .8269)N)$$

gives time for muon m_μ decay.

For π^0 decay time $m_v \rightarrow m_e$ (E&M decay) along with $8.87 \rightarrow 7 = m_p/m_\pi$ in the above equation.

For resonances $m_v \rightarrow m_e$ (E&M decay) in 9.23 gives time of decay.

Note the second term here contains a $i=-1$ and so it is a exponential decay term e^{-Et} with $.693/E=t$ the "half life".

Thus we get π^0 , $\pm\pi$, K mesons and hyperon, muon, neutron, resonance half lives from (these modifications of) equation 9.23.

9.7 CASE 2 Excited State F, charge=0. 2P1/2

Recall from 9.4 that case 1 implies $E_q \rightarrow m$ in case 2 (in 9.4). Also

$1/g_{rr} \approx 1 - k_H/k_H + \epsilon = \epsilon$ for $-\epsilon$. Net charge=Zero. Thus let $R=k_H+r$, $r < R$, $r'=k_H\epsilon+r$

$$\left[\left(\frac{dt}{ds} \sqrt{g_{00}} m \right) + m \right] F - \hbar c \left(\sqrt{g_{rr}} \frac{d}{dr} + \frac{j+3/2}{r} \right) f = 0$$

$$\left[\left(\frac{dt}{ds} \sqrt{g_{00}} m \right) - m \right] f + \hbar c \left(\sqrt{g_{rr}} \frac{d}{dr} - \frac{j-1/2}{r} \right) F = 0$$

Recall equations 8.1, 8.2:

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] F - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{j+3/2}{r} \right) f = 0$$

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] f + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{j-1/2}{r} \right) F = 0$$

$$\sqrt{\kappa_{rr}} = \frac{1}{\sqrt{1 - \frac{r_H}{R} + \varepsilon}} \approx \frac{\sqrt{R}}{\sqrt{R - r_H + R\varepsilon}} = \frac{\sqrt{r_H + r}}{\sqrt{r_H + r - r_H + (r_H + r)\varepsilon}} \approx$$

$$\frac{\sqrt{r_H + r}}{\sqrt{r_H \varepsilon + r(1 + \varepsilon)}} \approx \frac{\sqrt{r_H}}{\sqrt{r_H \varepsilon + r}} = \frac{\sqrt{r_H}}{\sqrt{r'}}$$

Also $\left(\frac{dt}{ds} \right) \sqrt{\kappa_{00}} \rightarrow E$ in the Dirac equation 18.1. Therefore equation 19.1 reads: $r' = r_H \varepsilon + r$

$$[E + m]F - \hbar c \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr} + \frac{j + \frac{3}{2}}{r_H + r} \right) f =$$

$$[E + m]F - \hbar c \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr} + \frac{j + \frac{3}{2}}{r_H + r} \right) f = 0$$

$$[E - m]F - \hbar c \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} + \left(1 - \frac{r}{r_H} \right) \frac{j + \frac{3}{2}}{r_H} \right) f = 0. \quad \text{and}$$

$$[E - m]f + \hbar c \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr} - \left(1 - \frac{r}{r_H} \right) \frac{j - \frac{1}{2}}{r_H} \right) F = 0 \quad \text{Thus}$$

.

$$f = -\frac{\hbar c}{(E - m)} \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} - \left(1 - \frac{r}{r_H} \right) \frac{(j - \frac{1}{2})}{r_H} \right) F \quad \text{Therefore}$$

$$[E + m]F - \hbar c \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} + \left(1 - \frac{r}{r_H} \right) \frac{j + \frac{3}{2}}{r_H} \right) f = 0 \quad \text{using } r = r' + r_H \varepsilon$$

$$(E + m)F - \hbar c \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} + \left(1 - \left(\frac{r' - \varepsilon r_H}{r_H} \right) \right) \frac{j + \frac{3}{2}}{r_H} \right) \frac{-\hbar c}{E - m} \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} - \left(1 - \frac{(r' - \varepsilon r_H)}{r_H} \right) \frac{j - \frac{1}{2}}{r_H} \right) F = 0$$

Multiplying both sides by $|E - m_p|$ we obtain:

$$\left(\frac{E^2 - m^2}{(\hbar c)^2} \right) F + \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} + \left(1 - \frac{(r' - \varepsilon r_H)}{r_H} \right) \left(\frac{j + \frac{3}{2}}{r_H} \right) \right) \left(\sqrt{\frac{r_H}{r'}} \frac{d}{dr'} - \left(1 - \frac{r' - r_H \varepsilon}{r_H} \right) \left(\frac{j - \frac{1}{2}}{r_H} \right) \right) F = 0$$

$$\frac{E^2 - m^2}{(\hbar c)^2} - \left((1 + 2\varepsilon) \left(\frac{(j + \frac{3}{2})(j - \frac{1}{2})}{r_H^2} \right) \right) F + \sqrt{\frac{r_H}{r'}} \left(\frac{j - \frac{1}{2}}{r_H^2} \right) + (1 + \varepsilon) \frac{(j + \frac{3}{2})}{r_H} \sqrt{\frac{r_H}{r'}} \frac{d}{dr'} +$$

$$\left(\frac{r_H}{r'} \frac{d^2}{dr'^2} - \frac{1}{2} \frac{r_H}{r'^{4/2}} \frac{d}{dr'} \right) F = 0 \quad \text{Multiplying both sides by } r'^2 \text{ we obtain:}$$

$$\left(\left[\frac{E^2 - m^2}{(\hbar c)^2} r'^2 \right] - (1 + 2\varepsilon) \left(j + \frac{3}{2} \right) \left(j - \frac{1}{2} \right) \left(\frac{r'^2}{r_H^2} \right) \right) F + (1 + \varepsilon) \left(\frac{j + \frac{3}{2}}{\sqrt{r_H}} \right) r'^{3/2} \frac{d}{dr'} +$$

$$\left[\left(r' r_H \frac{d^2}{dr'^2} \right) - \frac{1}{2} r_H \frac{d}{dr'} - \frac{\sqrt{r_H} r'^{3/2}}{r_H^2} \left(j - \frac{1}{2} \right) \right] F =$$

Defining $r' \equiv r^2$ and doing the derivatives in the new variable:

$$\frac{dF}{dr^2} = \frac{dF}{dr} \frac{dr}{dr^2} = \frac{1}{2r} \frac{dF}{dr} \quad \text{and}$$

$$\frac{d^2 F}{dr^2} = \frac{1}{2r} \frac{d}{dr} \left(\frac{1}{2r} \frac{dF}{dr} \right) = \frac{1}{2r} \left(-\frac{1}{2r^2} \frac{dF}{dr} \right) + \frac{1}{2r} \frac{1}{2r} \frac{d^2 F}{dr^2} =$$

$$-\frac{1}{4r^3} \frac{dF}{dr} + \frac{1}{4r^2} \frac{d^2 F}{dr^2} \quad \text{Substituting these expressions for the derivatives in:}$$

$$\left[\frac{E^2 - m^2}{(\hbar c)^2} r^4 - (1 + \varepsilon) \left(j + \frac{3}{2} \right) \left(j - \frac{1}{2} \right) \left(\frac{r^4}{r_H^2} \right) \right] F +$$

$$\left[\frac{r_H}{4} \frac{d^2}{dr'^2} - \frac{r_H}{4r} \frac{d}{dr'} \right] F - \frac{\sqrt{r_H} r'^3}{r_H^2} \left(j - \frac{1}{2} \right) F + \frac{r^3}{r_H} (1 + \varepsilon) \left(j + \frac{3}{2} \right) \left(\frac{\sqrt{r_H}}{2r} \right) \frac{d}{dr'} F - \frac{r_H}{4r} \frac{d}{dr} F =$$

$$\left[\left(\frac{E^2 - m^2}{(\hbar c)^2} \right) - \frac{(1 + 2\varepsilon) \left(j + \frac{3}{2} \right) \left(j - \frac{1}{2} \right)}{r_H^2} \right] \sum a_n r^{n+4} + \frac{r_H}{4} \sum (n - 1) n a_n r^{n-2} -$$

$$\frac{r_H}{4} \sum n a_n r^{n-2} - \frac{\sqrt{r_H}}{r_H^2} \left(j - \frac{1}{2} \right) \sum a_n r^{n+3} + (1 + \varepsilon) \frac{(j + \frac{3}{2})}{2\sqrt{r_H}} \sum n a_n r^{n+1} - \frac{r_H}{4} \sum n a_n r^{n-2} =$$

$$=$$

$$\left(\frac{E^2 - m^2}{(\hbar c)^2} - (1 + 2\varepsilon) \frac{(j + \frac{3}{2})(j - \frac{1}{2})}{r_H^2} \right) \sum a_{n-4} r^n + \frac{r_H}{4} \sum (n + 1)(n + 2) a_{n+2} r^n -$$

$$\frac{r_H}{4} \sum (n + 2) a_{n+2} r^n - \frac{\sqrt{r_H}}{r_H^2} \left(j - \frac{1}{2} \right) \sum a_{n-3} r^n + (1 + \varepsilon) \frac{(j + \frac{3}{2})}{2\sqrt{r_H}} \sum (n + 1) a_{n-2} r^n -$$

$$\frac{r_H}{4} \sum (n + 2) a_{n+2} r^n = 0$$

Combining terms noting simplification due to combining the a_{n+2} terms

$$\left(\frac{E^2 - m^2}{(\hbar c)^2} - (1 + 2\varepsilon) \frac{(j + \frac{3}{2})(j - \frac{1}{2})}{r_H^2} \right) \sum a_{n-4} r^n + \frac{r_H}{4} \sum (n - 1)(n + 1) a_{n+2} r^n +$$

$$\frac{\sqrt{r_H}}{r_H^2} \left(j - \frac{1}{2} \right) \sum a_{n-3} r^n + (1 + \varepsilon) \frac{\left(j + \frac{3}{2} \right)}{2\sqrt{r_H}} \sum (n-1) a_{n-1} r^n = 0. \quad ?$$

Next we write the individual eigenfunctions as:

$$\left(\frac{E^2 - m^2}{(\hbar c)^2} - (1 + 2\varepsilon) \frac{\left(j + \frac{3}{2} \right) \left(j - \frac{1}{2} \right)}{r_H^2} \right) \sum a_{n-4} r^n = 0$$

Thus since these series terms add to zero:

$$E = \sqrt{m^2 + (\hbar c)^2 (1 + 2\varepsilon) \frac{\left(j + \frac{3}{2} \right) \left(j - \frac{1}{2} \right)}{r_H^2}} \quad (9.24)$$

$$(1 + \varepsilon) \frac{j + \frac{3}{2}}{2\sqrt{r_H}} \sum (n-1) a_{n-2} r^n = 0 \quad \text{Here } r' = r^2 \text{ so } r^{-2/2} = r'^{-1} = \sqrt{r}^{-2}$$

$$\frac{r_H}{4} \sum (n^2 - 1) a_{n+2} r^n = 0 \quad (9.25)$$

-

$$-\frac{\sqrt{r_H}}{r_H^2} \left(j - \frac{1}{2} \right) \sum a_{n-3} r^n = 0. \quad (9.26)$$

J=1/2 with N=1 solves the indicial equation implied by 9.24-9.26. Recall from 9.4 that m=proton in this case (case 2). The energy in 9.24 is then that of a neutral particle (q=0) with the mass of the neutron so $E = E_q = m = m_N$. See equation 9.23b for neutron lifetime and 2P3/2 for neutron spherical harmonic state, section 10.3) But in case 2 and equation 9.23 then the previously derived charged spin 1/2 hadrons m_Σ , m_Ξ can also be put back into the Dirac equations for 'm' (instead of the proton). Thus the charged, m_Σ , m_Ξ from equation 9.14 can be put into the "m" in 9.24 which gives the **neutral** $E = m = m_N$, m_Ξ . m_Σ has a N=1/2 and so does not satisfy the above equations and so does not exhibit a stable neutral Σ . Recall the Ω^- (which is J=3/2) is not J=1/2 so doesn't have a neutral counterpart as does the proton and these other J=1/2 hyperons.

Recall the iterated Dirac equation is the Klein Gordon (in χ with J=0) equation eigenstate transitions.

J=0, q=0 Case 2

Recall J=0 is allowed in every case.

m=1 proton, j=0 in equation 9.24 means K Long. Equation 9.23 gives K long mass eigenvalue:

$1 + (0 + 3/2)(0 - 1/2)/1 = 1/4$. Thus $\sqrt{.25} = .5$. Thus **.5X938X1.06=497 MeV=K_{long}**. Note case 2 is zero charge and note also from section 9.8 that the Strangeness=2| $\sqrt{.5}$ |=2*.707 ≈ 1 as in strangeness equation 9.34 below.

$m \approx 1$ for Neutron then in 9.24 we have K short, if $m = m_\Xi$ and J=0 then D° Long.

If $m = m_\Xi$ j=0, and neutral then 9.24 gives D° Short.

9.8 CASE 3 m=0, so ψ_L , f state, charge=0 (lower case of equation 9.5).

In case 3 there is no central force therefore N=0 and j=1/2 in f. This is the m=0 left handed doublet case of Chapter 3. Let $R = k_H - r$, $r \ll R$ for stability we can write:

$$\sqrt{K_{rr}} = \frac{1}{\sqrt{1+\frac{r_H}{R}+\varepsilon}} \approx \frac{\sqrt{R}}{\sqrt{R+r_H+R\varepsilon}} =$$

$$\frac{\sqrt{r_H-r}}{\sqrt{r_H-r+r_H+(r_H-r)\varepsilon}} =$$

$$\frac{\sqrt{r_H-r}}{\sqrt{r_H(2+\varepsilon)-r(1+\varepsilon)}} =$$

$$\frac{\left(1-\frac{\varepsilon}{4}\right)}{\sqrt{2}} \left[\frac{1-\frac{r}{2r_H}+\frac{r^2}{8r_H^2}+..}{1-\frac{r}{4r_H}+\frac{r^2}{16r_H^2}+..} \right]$$

$$\left(\frac{1-\frac{\varepsilon}{4}}{\sqrt{2}}\right) \left(1 - \frac{r}{4r_H} + \frac{3r^2}{32r_H^2} - ..\right) \approx \frac{1-\frac{r}{4r_H}}{\sqrt{2}}$$

Therefore equation 9.1 reads:

$$[E + m_p]F - \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{j+\frac{3}{2}}{r_H-r} \right) f = 0$$

$$[E - m_p]F - \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{\left(1 + \frac{r}{r_H}\right)\left(j + \frac{3}{2}\right)}{r_H} \right) f = 0$$

$$[E - m_p]f + \hbar c \left(\sqrt{K_{rr}} \frac{d}{dr} - \frac{j-\frac{1}{2}}{r} \right) F = 0 \quad (9.27)$$

From the above equation 9.27 if (and $j=\frac{1}{2}$) $m_p=0$ then

$$[E - m_p]f + \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} - \frac{\left(1+\frac{r}{r_H}\right)\left(j-\frac{1}{2}\right)}{r_H} \right) F = 0$$

Therefore (with $j=\frac{1}{2}$) from equation 9.27 for small r. In any case:

$$F = \hbar c \frac{\hbar c}{(E+m_p)} \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{\left(1+\frac{r}{r_H}\right)\left(j+\frac{3}{2}\right)}{r_H} \right) f$$

$$[E + m_p]F - \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{j+\frac{3}{2}}{r_H-r} \right) f = 0$$

Solving for f and substituting back in 9.27

•

$$\begin{aligned}
& [E - m_p]f + \hbar c \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} - \frac{\left(1 + \frac{r}{r_H}\right)\left(j - \frac{1}{2}\right)}{r_H} \right) \\
& \frac{\hbar c}{E + m_p} \left(\left(1 - \frac{r}{4r_H}\right) \frac{d}{\sqrt{2}dr} + \frac{\left(1 + \frac{r}{r_H}\right)\left(j + \frac{3}{2}\right)}{r_H} \right) F = [E - m_p]f + \\
& \frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \left(-\left(1 - \frac{r}{4r_H}\right) \frac{d}{r_H 4\sqrt{2}dr} + \left(1 - \frac{r}{4r_H}\right)^2 \frac{d^2}{\sqrt{2}dr^2} + \frac{\left(1 - \frac{r}{4r_H}\right)\left(j + \frac{3}{2}\right)}{r_H^2} \right) f \\
& + \frac{(\hbar c)^2}{E + m_p} \left(-\left(1 + \frac{3r}{4r_H}\right)\left(j - \frac{1}{2}\right) \frac{d}{\sqrt{2}r_H dr} - \frac{\left(1 + \frac{r}{r_H}\right)^2 \left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^2} \right) f = \\
& \left([E - m_p] + \left[\frac{(\hbar c)^2}{(E + m_p)} \left(-\frac{\left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^2} + \frac{\left(j + \frac{3}{2}\right)}{\sqrt{2}r_H^2} \right) \right] \right) f + \\
& \frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \left(\frac{2\sqrt{2}\left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^2} - \frac{\left(j + \frac{3}{2}\right)}{4r_H^2} \right) rf + \\
& \frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \left(-\frac{1}{k_H 4\sqrt{2}} + \left(j - \frac{1}{2}\right) \frac{1}{k_H} \right) \frac{df}{dr} + \\
& \frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \left(-\frac{1}{4\sqrt{2}r_H} + \frac{\left(j - \frac{1}{2}\right)}{r_H} \right) \frac{df}{dr} + \\
& \left(\frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \right) \left(\frac{1}{16\sqrt{2}r_H^2} - \frac{3\left(j - \frac{1}{2}\right)}{4r_H^2} \right) r \frac{df}{dr} + \\
& \left(\frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \right) \left(\frac{1}{\sqrt{2}} \right) \frac{d^2 f}{dr^2} + \\
& \left(\frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \right) \left(\frac{-1}{2\sqrt{2}r_H} \right) r \frac{d^2 f}{dr^2}
\end{aligned}$$

Next substitute in $F = \sum_n a_n r^n$

$$\begin{aligned}
& \sum_M^N \left([E - m_p] + \left[\frac{(\hbar c)^2}{(E + m_p)} \left(-\frac{\left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^2} + \frac{\left(j + \frac{3}{2}\right)}{\sqrt{2}r_H^2} \right) \right] \right) a_n r^n + \\
& \sum_M^N \frac{(\hbar c)^2}{(E + m_p)\sqrt{2}} \left(\frac{2\sqrt{2}\left(j + \frac{3}{2}\right)\left(j - \frac{1}{2}\right)}{r_H^3} - \frac{\left(j + \frac{3}{2}\right)}{4r_H^3} \right) a_{n-1} r^n + (9.28)
\end{aligned}$$

$$\sum_M^N \left(\frac{(\hbar c)^2}{(E+m_p)\sqrt{2}} \right) \left(-\frac{1}{4\sqrt{2}r_H} + \frac{(j-\frac{1}{2})}{r_H} \right) (n+1)a_{n+1}r^n \quad (9.29)$$

$$\sum_M^N \left(\frac{(\hbar c)^2}{(E+m_p)\sqrt{2}} \right) \left(\frac{1}{16\sqrt{2}r_H^2} - \frac{3(j-\frac{1}{2})}{4r_H^2} \right) na_n r^n + \quad (9.30)$$

$$\sum_M^N \left(\frac{(\hbar c)^2}{(E+m_p)\sqrt{2}} \right) \left(\frac{1}{\sqrt{2}} \right) (n+2)(n+1)a_{n+2}r^n + \quad (9.31)$$

$$\sum_M^N \left(\frac{(\hbar c)^2}{(E+m_p)\sqrt{2}} \right) \left(\frac{-1}{2\sqrt{2}r_H} \right) (n+1)na_{n+1}r^n = 0 \quad (9.32)$$

We now take 9.27 and 9.30 together and 9.29 and 9.32 together (since they have the same a_n). Thus there are 4 independent series (with 9.28 and 10.31) here. The equation 9.27 and 9.30 nth terms give:

$$[E + m_p] + \left[\frac{(hc)^2}{E - m_p} \left(-\frac{(j + \frac{3}{2})(j - \frac{1}{2})}{r_H^2} + \frac{(j + \frac{3}{2})}{\sqrt{2}r_H^2} \right) \right] +$$

$$\left(\frac{(hc)^2}{(E - m_p)\sqrt{2}} \right) \left(\frac{1}{16\sqrt{2}r_H^2} - \frac{3(j - \frac{1}{2})}{4r_H^2} \right) n.$$

At some value of $n=N$ we have for a solution

$$(E^2 - m_p^2) + \left(-\frac{(j+\frac{3}{2})(j-\frac{1}{2})}{r_H^2} + \frac{(j+\frac{3}{2})}{\sqrt{2}r_H^2} \right) + \frac{1}{\sqrt{2}} \left(\frac{1}{16\sqrt{2}r_H^2} - \frac{3(j-\frac{1}{2})}{4r_H^2} \right) N = 0 \quad \text{therefore rearranging:}$$

$$E = \sqrt{m_p^2 + \frac{1}{r_H^2} \left((j + 3/2)(j - 1/2) + .7071j + 1.0607 + (.0156 - (j - .5).5303)N \right)} \quad (9.33)$$

Recall from the equation 9.4 ‘f’ case that we have $m_p=m=0$, and zero charge therefore no central force thus $N=0$ in $f \propto r^0$ in equation 8.1. Therefore since there is small r and $dr^0/dt=d1/dt=0$ in the equations just above equation 9.27 along with 9.33 then the 9.27-9.32 equations add to zero and thus are solved. Also the $j=3/2$ (so $L=1$) case is not allowed since that requires a central force to give $L \neq 0$, $j=1/2$ and of course $j=0$ is allowed here. Thus

$N=0$, $j=1/2$, $m=0$ then from 9.33 we have $E=1115.8 \text{ Mev } \Lambda_0$

$N=0$, $j=0$, η mass and also gives $m=.56$ (with $m=0$) in 9.33 used in gyromagnetic ratio calculation for f . Recall $\varepsilon=.08$ (with $m=0$) for F in 9.14. This is the nontrivial f state zero point energy for $r < k$ since $\Psi=\psi+\chi$ from our observability definition. Note Kaons then give no strange bound states because this mass is real (in contrast to the imaginary pion mass in 9.22).

9.9 Strangeness

Recall that in 9.14 (which applies to Case 1 and Case 2) the energy is $E^2=m_0^2 + (j^2+1.7071j-1.10355-(j(.53033))+.7642)N)/k_H^2$. Now m_0^2 and E is conserved (m_0 is a constant) here and thus it appears that energy conservation implies that the square root of $j^2+1.7071j-1.10355-(j(.53033)+.7642)N \equiv S$ must be conserved. Therefore $E^2=m^2 + S^2$ then and “ S ” is conserved for the charged core states and thus for the neutrals given that in section 9.8 that $E_q \rightarrow m$ then (for f state $m=0$ we also have $S \approx E$ for Λ). We could also write $E^2=m^2 + C^2$ for the next $2P$ state eigenstates (call C charm if you want) which would also have their own associated production (since $\langle | \rangle$ not zero). Thus, as an example, normalizing to a factor of $2X$:

$$\begin{aligned}
2XSQR[(.5(.5303)+.7642)(0)] &= 0 = S_{\text{nucleon}}, & 2XSQR[(.5(.5303)+.7642)(-1)] &\approx 2 = S_{\Xi}, \\
2XSQR[(.5(.5303)+.7642)(-1/2)] &\approx 1 = S_{\Sigma}, & 2XSQR[(1.5^2+1.7071(1.5)-1.10355- \\
&& (1.5(.5303+.7642))(-1)] &\approx 3 = S_{\Omega}.
\end{aligned}
\tag{9.34}$$

Strangeness is only an approximate conservation law in the examples in 9.34 but there is enough conservation at least for the “associated production” and we have not yet included the weak interaction here. This is a **direct derivation of strangeness**, instead of just having postulated it as it is in the standard model and QCD. Strangeness isn’t strange anymore.

Charm, bottom, top: In chapter 9 equation assuming hard spherical shell. We obtain other (less stable, resonances) particle groups using equation 9.5 by taking the quadratic approximation of g_{rr} (i.e., include the $(3/32)(r/k_H)^2$ term in 9.5) Using 10.8 instead of just the linear approximation we used above. Recall that the perturbative $(3/32)(r/k_H)^2$ term had to be included since it gave a ≈ 20 MeV correction to the hyperon masses.

C Meson Mass Derivation From Potential Of Chapter 10 And The New Pde eq.9

C Spherically Symmetric Wave Function Required

```

PROGRAM FracsN
DOUBLE PRECISION A,B,C,D,E,F,H,I,I1,J,KK
DOUBLE PRECISION K1,K2,K3,K4,N1,N2,N3,N4,R,W,X,Y,Z
DOUBLE PRECISION Y1,E1,E2,MM1,MM2,MM3,EE,JJ
integer N,M,M1
DIMENSION EE(400)
C Variational principle on E with respect to I and Y1,
C RungeKutte on D equation 8.1. Y=2 width Deuteron
C pion oscillation resonance modeled between 0 and Y=2.
H=0.001
mH=2 !harmonic number for oscillation inside Y=2.
C mN=1 gives pion 0 and K+-,mN=2 gives pi+- and Ko resonance
ep=0.08*mH !pion 1st and 2nd harmonic resonance added to Y1
W=1.0+ep !pion mass added to nucleon.
J=0.0 !spin 0 mesons
X=0.0001 !mass energy increments
I1=100000000.0
A=0.0
B=0.0
C=0.0
E=0.0
KK=78.8 !gives MeV energy units
JJ=J*1.
Y1=2.0+ep !pion increases Y1.
50 D=.00000001
I1=0.0
F=.00000001
Y=Y1
60 R=Y
V=1.0/(1.0+ep-R) !chapter 14 potential for spin 0
E1=E

```

```

K1=((W-E-V)*F)+(((J-0.5)/R)*D)
N1=((E+W+V)*D)-(((J+1.5)/R)*F)
R=R+(0.5*H)
V=1.0/(1.0+ep-R)
K2=((W-E-V)*(F+(0.5*H*N1)))+(((J-0.5)/R)*(D+(0.5*H*K1)))
N2=((E+W+V)*(D+(0.5*H*K1)))-(((J+1.5)/R)*(F+(0.5*H*N1)))
K3=((W-E-V)*(F+(0.5*H*N2)))+(((J-0.5)/R)*(F+(0.5*H*K2)))
N3=((E+W+V)*(D+(0.5*H*K2)))-(((J+1.5)/R)*(F+(0.5*H*N2)))
R=R+(.5*H)
V=1.0/(1.0+ep-R)
K4=((W-E-V)*(F+(H*N3)))+(((J-0.5)/R)*(D+(H*K3)))
N4=((E+W+V)*(D+(H*K3)))-(((J+1.5)/R)*(F+(H*N3)))
E=E1
F=F+((H/6.0)*(N1+(2.0*N2)+(2.0*N3)+N4))
D=D+((H/6.0)*(K1+(2.0*K2)+(2.0*K3)+K4))
I=(F*F)+(D*D)
100 I1=I1+(I*(R+(0.5*H))*(R+(0.5*H)))
    IF((abs(R-1.0-ep)).LT.(0.9*H))THEN
        Y=Y-(2.0*H)
        GOTO 60
    ENDIF
    Y=Y-H
    IF(Y.LT.0.0)THEN
        GOTO 200
    ENDIF
    GOTO 60
200 E=E+X
    C=I1
    IF(B.LT.A)THEN

        GOTO 310
    ENDIF
    GOTO 312
310 IF(C.GT.B)THEN
    ENDIF
312 IF(B.GT.A)THEN
    GOTO 315
    ENDIF
    GOTO 320
315 IF(C.LT.B)THEN
    print *, ' '
    print *, 'E=',(E-X)*KK, ' J=',J, ' max I'
    ENDIF
320 IF(E.GT.8.0)THEN
    GOTO 349
    ENDIF

```

```

      A=B
      B=C
330  GOTO 50
349  print*,'program finished'
350  stop
      End

```

C Results for spin 0, L=0 are

C For $m_N=1$ get 135 MeV π^0 and 493 K^\pm for resonance with 1 meson.

C For $m_N=2$ get 139 MeV π^\pm and 497 MeV K^0 for resonance with two

497 MeV K^0 for resonance with two mesons in ordinary nuclear matter nucleus would split before K energy created. In a neutron star however K's could be created.

This Fortran computer program only requires a few seconds to run on a PC. On the other hand lattice gauge theory programs (assuming a SU(3) lattice) require massive computing power and really do not duplicate high energy liquid state strong interactions anyway.

Here the pion is a $r=2R_H$ proton with no net rotation and the central electron in a $m=0$ state so net spin = 0. $\sigma \gg 1/20$ Barn so annihilation occurs outside r_H and the pion decays.

Stability of 3e Composite

The positrons are ultrarelativistic, $\gamma=917$, so the field lines are Fitzgerald contracted so so the E&M field lines are contracted resulting in our explanation of the strong force.

$dt'^2 = \kappa_{00} dt^2 = (1 - r_H/r) dt^2$ is zero if $r=r_H$ so clocks stop so stability. But some ψ must leak out so some electron positron annihilation must occur. But as we show below virtual decay and annihilation still results in stability




Fig3

So this is a **virtual annihilation-creation process** inside r_H , **implying that this two positron-single electron state is stable** (yet another reason for baryon stability). See eq.11 also. We rigorously derive the low mass ($<3\text{GeV}$) hyperon eigenvalues using the Frobenius series solutions to eq.11 near $r=r_H$ (from 3rdpt, $r \approx r_H$) in Ch.8-Ch.11.

B field Inside r_H

That above eq.1 calculation used $r_H = e^2/m_e c^2$, $q/t=i$. $q=e=1.6 \times 10^{-19}$ C, $u_0=4\pi \times 10^{-7}$, $A=\pi r_H^2$, $BA=2 \times 10^{-15} \text{Wb}$, $BA = \left(\frac{\mu_0 i}{2r_H} \right) \pi r_H^2$. This allows us to calculate $B=10^{11} \text{T}$ inside r_H . Note this creation and annihilation of this B field (above fig/3) must imply a Faraday's law Meisner effect zero point energy $J=0$ eq.9.23 pion cloud meaning there will be a constant cloud of moving pions around baryons, thereby proving the existence of the well known pion caused Yukawa force. (potential $V=e^{-kr}/r$). One interesting consequence is that just after a typeII supernova the protons are most compacted (reaction force to the outward action push) and so the pion cloud is squeezed out. So there is no Mesiner effect and so the entire neutron star exhibits this huge 10^{11}T B field (thereby explaining magnetars) But the star eventually radiates this (potential) energy away expanding the star and so not squeezing out the pions anymore and so the Meisner effect suddenly returns thereby with a sudden change of B flux creating a large Faradays law EMF thereby explaining FRBs (energy= $(B^2/(2u_0))X[(4/3)\pi(10^4)^3]=(10^{11})^2/(2 \times 4\pi \times 10^{-7}) \times 4 \times 10^{12}=1.4 \times 10^{40} \text{Joules}$ FRB enough energy for a ms pulse to be detected anywhere in the (10^{12}Ly radius) universe even by modest Jansky sensitivity RF receivers..).

Chapter 10

$r \approx r_H$ Application: $2P_{3/2}$ Half Integer Spherical Harmonics Solutions. This is a continuation of Chapter 9

10.2 Overview of $2P_{3/2}$ Solutions to Equation 9 (the New Dirac equation) at $r \approx r_H$ in the Context of the Equivalence Principle (single charge e) Implication


Allowing this *single* charge 'e' to move near and inside that stable singularity radius $r \approx 2q^2/mc^2$ in the $\sqrt{g_{ij}}$ in this new Dirac equation (equation 2) as we see below makes the motion relativistic but stable requiring all the Dirac equation spherical harmonic solutions, not just the ones allowed by the Schrodinger equation. Also the next order of approximation above the hard shell for our g_{00} horizon $r_H = 2e^2/m_e c^2$ is the harmonic oscillator $V \propto r^{+1}$ giving the $SU(3)$ SYMMETRY of the three dimensional harmonic oscillator. The $+1$ in the exponent of V (instead of the inverse square law-1) also reverses the sign on the exchange integral $\pm \int \psi_{111}^*(r') \psi_{lmn}^*(r'') V(r', r'') \psi_{lmn}(r') \psi_{111}(r'') d\tau = J$ designating the symmetric and antisymmetric states), making here then the $J=3/2$ state $m=-3/2$ and $3/2$

(i.e., $\psi = \mathcal{Y}^{3/2}_{3/2}(\theta, \phi) + \mathcal{Y}^{-3/2}_{3/2}(\theta, \phi) = 2P_{3/2}$ eigenspinor) the first ground state that varies with azimuthal angle (baryons) above the already filled 1S (in analogy with helium) on the energy ladder instead of the expected $1/2$ and $-1/2$ (these $1/2$ s by the way give $2P_{1/2}$ in the $\psi^* \psi$ of the next higher P orbital slots) that vary with azimuthal angle (baryons).

Also recall the identity $(\exp(i\phi) + \exp(-i\phi))/2 = \cos\phi$. The $\mathcal{Y}^{3/2}_{3/2}$ orbital is a $\exp(i3/2\phi)$ and $\mathcal{Y}^{-3/2}_{3/2}$ orbital is $\exp(-i3/2\phi)$ and thus from the identity the summed state is $\cos(3/2\phi)$ with probability density $\psi^* \psi = \cos^2(3/2\phi)$, the trifolium three lobed shape. Thus there are TWO +e s giving a net charge of $+2/3e$ in each lobe because the +electron charge 'e' is in each orbital lobe on the average only $1/3$ the time (FRACTIONAL CHARGE) giving the many scattering properties (such as jets) associated with the angular distribution of multiple fractional charges interior to this horizon. The lobe 'structure' *can't leave* (ASYMPTOTIC FREEDOM) as in the Schrodinger equation case or move so is NONRELATIVISTIC in contrast to its rapidly moving m_e constituent.

Finally we solve the problem with the new pde using a computer program, set the boundary conditions as if the Deuteron was a square well. See end of chapter 9 for the fortran program. In any case we can build the hyperons and mesons with integer charges e, don't need the fractional charges.

10.3 Trifolium Diagram



2P_{3/2} solutions to Dirac equation. $\sqrt{\kappa_{\mu\mu}} \gamma^\mu \frac{\partial \psi}{\partial x^\mu} - \omega \psi = 0$ $\kappa_{00} = 1 - \frac{r_H}{r}$ Stability at $r \approx r_H$ since then $\kappa_{00} = 0$

Ultrarelativistic LS coupling fills 2P_{3/2} first $\kappa_{00} = \frac{1}{\kappa_{rr}}$

Exchange integral implies (after filled 1S +2e) next bound state =

$$\psi = \mathcal{Y}_{3/2}^{3/2} + \mathcal{Y}_{3/2}^{-3/2} = \frac{e^{i\frac{3}{2}\phi} + e^{-i\frac{3}{2}\phi}}{2} = \cos\left(\frac{3}{2}\phi\right)$$

$$\psi^* \psi = \cos^2\left(\frac{3}{2}\phi\right) = \text{trifolium}$$

Electron charge e spends 1/3 of its time in each lobe making each lobe (1/3)e charged on average. For two such electrons it is (2/3)e.

Start with 1S state filled singlet = (1/2 -1/2) = γ_0^0 . Add 2P_{3/2} and electron -e for bound state e.g., proton. Add to this proton 2P_{3/2} state -e and +e in filled 2P_{1/2} and a third -e with spin down to get neutron (spin 1/2). Singlet = π^0 with one of the electrons $\Delta\epsilon$ in the first electron excited state ϵ (muon). Add -e and +e in filled 2P_{1/2} and a third +e with down spin get π^+ (spin 0)



Figure 10-1 Trifolium diagram

$2P_{3/2}$ fills first in Aufbau principle for ultrarelativistic hard shell (Alfredo 1998).

Electron in limaçon lobe added to trifolium lobe to give bound state:

$-e + (2/3)e = -(1/3)e$ d. Add other two lobes $(2/3)e$ u + $(2/3)e$ u = uud =

Fill in rest of P states same way.



6 P orbital slots at $r=r_H$ Fill states as nondegenerate energy (level) goes up \longrightarrow

Possible SHM interaction between these lobes gives excited states.

LS coupling Lande' g-factor structure gives minimal LS energy for smallest L

So net spin 1/2 states preferred.

Fig2 Note we get a similar shape to the trifolium with lattice QCD theory.

Fig4





Fig.3 Second diagram is the E field of a ultrarelativistic charged particle ("Electronic Motion", McGraw Hill, Harman)

Thus the neutron charge configuration allows for the creation of both the W and the H_v and the proton charge configuration does not allow this.



$2P_{3/2}$ at $r=r_H$
 positive 1e charge
 $J_A + J_C + J_B =$
 $(-.5+1)+(-.5+1)-.5=.5$ spin
 ultrarelativistic electron motion gives
 correct mass.
 plate field from object A does not intersect
 particle C so bound state.

$2P_{3/2} + 2P_{1/2}$
 D not bound so He
 $H_v + H_e = 0$
 So left handed neutrino
 exceeds epsilon so W S
 Matrix

Fig5

10.5 Deuteron is the $J=1, \psi \propto r \sin \theta \propto Y^1_1(\theta, \phi)$ (at $r=r_H$) solution of the Newpde.eq.9.14

It is then a double lobed $\psi^* \psi$ along the z axis at $r=r_H$. Deuteron state is derived from the above Ch.9 above Frobenius series solution to the Newpde. From equation 9.14 we find with these inputs that

$$E = 1867 \text{ Mev} \quad (9.23)$$

Lorentz contraction
 $z=1$
 $2\text{bonds} = B\#$
 $\frac{e^2}{r_H^2}$
 $\frac{e^2}{r_H^2}$
 $wobble$

Equation 9.23 $N=-1, J=1 Y^1_1(\theta, \phi)$ solution to the NewPde at $r=r_H$

$PE = \frac{e^2}{r_H} = 1.06 \text{ Mev} = \frac{1}{2} k r_H^2 = \hbar \omega (n + \frac{1}{2}) = \text{SHM} = \text{simple harmonic motion energy } z=0$
 $r_H = 2.8 \times 10^{-15} \text{ m}$
 $\frac{1}{2} m/2$ Circular motion Fitzgerald contraction from side
 Equipartition Of Energy $N = \text{SHM} = \text{LS} = \text{Angular velocity energy}$
 Diatomic degrees of freedom = 6. 3 translational plus 2 rotational
 don't count in binding energy leaving $f = 6 - (3+2) = 1$
 $(B\#)PE = BE$
 $(2)1.06 = BE$
 $2 \times 1.06 = 2.12 \text{ Mev} = \text{Deuteron Binding energy.}$

The potential energy between the central electron (for $z=1$) and the two $+e$ is $PE_e = e^2/r_H$
 $= 9 \times 10^9 (1.602 \times 10^{-19})^2 / (1.4 \times 10^{-15}) = 1.7 \times 10^{-13} \text{ J}$. (Tabulated D charge radius = $2.128 \times 10^{-15} \text{ m}$)
 $1.7 \times 10^{-13} / 1.602 \times 10^{-19} = 1.06 \times 10^6 \text{ eV} = 1.06 \text{ Mev} = B$ at $r=r_H, z=0$ side view

$(B\#)B = BE$ = binding energy

Deuterium D (2) $1.06 = 2.2 \text{ Mev}$

For many D (>3 rotating harmonic oscillators: Deuterons) use 3D harmonic oscillator as a first approximation.

Neutron

With the second proton missing a single electron has .78Mev instead of our .5 Mev binding energy (Neutron β decays hint at this number also.). The extra neutron mass energy beyond the proton ($2P_{1/2}$) \sim **half the Deuteron BE** $=5SHM/2 = 2.2/2 = 1.1$. For single bond $\approx .78$ **plus the electron** where $2e^2/r_H = SHM$. So half is $.78 + .511 = 1.3$ Mev the excess mass of a neutron above a proton.

Summary

The shell model sect.6.12 also requires this average radius and so has that inner and outer metric (sect.6.11, 3.1) as in sect 3.1 just like all solutions of

We actually understand the nucleus of the atom from first principles eq.9.23 Y_1^1 at $r=r_H$) $N=-1$, $J=1$ solution to the Newpde here. We even got the Deuteron binding energy this way.

The shell model can now be understood with each Deuteron having the same SHM excited states as every other deuteron in the nucleus given the equipartition of energy (recall the shell model mysteriously uses the excited state of just one SHM oscillator in a vacuum. But there are in general many oscillators here and the nucleus is not a vacuum!). **So we finally understand why the shell model works.** For a history of the alternative Shell model, also see study by A.E.S. Green in 1956).

Comparison with QCD SU(3)

Instead of the shell model one dimensional SHM rotating (that LS) we could use the equivalent 3D oscillator and directly transform between x,y,z oscillator modes. In that regard the 3D components of the above SHM tensor $A_{ij} = (1/(2m))(p_i p_j + m^2 \omega^2 x_i x_j)$ and components of L satisfy Poisson bracket relations SU(3). So by including as a perturbation the rotation, the 3D SHM version gives SU(3) symmetry in the S derived from those A_{ij} (Herbert Goldstein, 'Classical Mechanics' 2nd edition, pp.425) which holds in both the classical and QM case then. In that latter case SU(3) is gauged in by starting it out as a infinitesimal rotation, taking the limit to get it into the exponent (as in appendix A). So we have also just explained the *origin* of the adhoc and convoluted QCD SU(3) gauged alternative to that correct Frobenius series $N=-1, J=1, Y_1^1, r=r_H$ solution to nuclear physics.

10.8 High (>100Gev) Energy Solutions to the Newpde

Note at high energy the electrons in the $2P_{3/2}$ lobes (e.g., udd) would appear stationary, not averaged blob (density distributions) anymore. We are back to having single 'e' (not fractional "e") scatterers again. Thus at very high energies (>100GeV) single e (not fractional charge) should once again dominate scattering and we should no longer see these "jets" (which in the above context is mere P wave scattering) caused by higher probability emission in these trifolium lobe directions (QCD no longer gives correct answers because of this). Also note that r_H in κ_{00} is a hard shell and therefore Van der Waals type liquid equation of state at >100Gev energies. Note by the way that the 6th 2P resonance is observable at these energies.

Let $\langle A' |$ represent the outgoing scattering wave immediately after a incident plane wave scatters off V. Let $|A\rangle$ be the $2P_{3/2}$ hyperon state for $r=r_H$ having the V. Thus at $r=r_H$ V itself will have the $2P_{3/2} * 2P_{3/2} = \psi * \psi$ trifolium shape and thus commute with $|A\rangle$ since they constitute the same structure ($2P_{3/2}$ commutes with itself). So since V commutes with $|A\rangle$ **then $\langle A' |$ also is a $2P_{3/2}$ state** or we have $\langle A' | V | A \rangle = 0$ and so no scattering into such states. Thus a type of 'P wave scattering' results from an incident plane wave. Thus we explain the origin of the 'jets' that are otherwise ascribed to scattering off quarks.

Note that when the mean free path d during the interaction time is very short ($d \ll (1/3) 2\pi r_H$) there is no more smearing between the $2P_{3/2}$ lobes and we have scattering off of independent point particles and the $2P_{3/2}$ state ceases to be relevant in the scattering and so the jets disappear.

(jet quenching). Thus at extremely high energy the scattering is from charge e (not $1/3e$) again and there are no more jets above top energy. LEP actually observed this effect just before it was shut down.

10.9 Charge Independence Of The Strong Interaction

It is well known that the strong interaction is approximately the same magnitude between Neutron-Proton, Neutron-Neutron and Proton-Proton pairs and thus is 'charge independent'. Also note our theory deals with electrons only which only has charge dependence if certain QM effects are ignored. But recall the orthogonality of S and P states as in $\langle S|P \rangle = 0$, $\langle S|S \rangle = 1$, $\langle P|P \rangle = 1$ given all the superscript and subscript substates (e.g., S and m) are the same as well in the bra and kets. The ordinary nuclear interaction here is due to a covalent bond (sharing electrons) which is also a very strong interaction (bond) at $r=r_H$ and is dependent on the spin S and m state and not so much on the sign of the charge. Thus these QM (valence, spin) effects are very strong at $r=r_H$. Thus the charge independence of the strong interaction is really an S state independence and $2P_{3/2}$ state dependence at $r=r_H$ of a $2P_{3/2}$ structure interacting with an S state.




Fig.5

There are no gauges required in this theory and the QCD SU(3) is such a gauge. We have found that hadrons are excited states composed of these half integer spherical harmonic lobes.

Chapter 11 Scattering Cross-Sections

From the energy component of polarized representation of equation 8.1, 8.2: and using iterated (as in bosonic) section 9.13 $E^2 = p^2 c^2 + m^2 c^4 =$

$$E^2 = \left(\frac{1}{\sqrt{\kappa_{oo}}} \right)^2 = \frac{1}{1 - \frac{r_H}{r}} = \frac{r}{r - r_H} = \frac{r}{r - r_H} - \frac{r - r_H}{r - r_H} + 1 = \frac{r_H}{r - r_H} + 1 = V + k \quad (10.8.1)$$

Note the resemblance of $E^2 = p^2 c^2 + m^2 c^4$ to the Schrodinger equation if the $E^2 = k^2 + 1/(r - r_H) + 1$ of equation 10.8.1 is substituted into it. We interpret this equation as representing a bounded volume with energy $E = V + k$ therefore allowing us to use that V in the usual Gauge theory method and so substitute it into the ordinary Dirac equation as gauge force. term. So we use $1/(r - r_H)$ instead of $1/r$ in the Dirac equation S matrix.

We use the equation 4.1 source and proceed in the usual way of Bjorken and Drell (here $1/r \rightarrow 1/(r - r_H/2)$) to construct the one vertex S matrix for the new Dirac equation 9. Recall the $1/2$ came from the square root in equation 4.1. Thus the k in the integrand denominator is found from the result of our $V = -1/(r - r_H/2)$ potential in equation 10.8.1 instead of the usual Coulomb potential $1/r$ in the large r limit (so a free electron otherwise):

$$S_{if} \equiv iZ \frac{1}{V} \sqrt{\frac{m^2}{E_f E_i}} \bar{u}(p_f, s_f) \gamma^0 u(p_i, s_i) \int_0^\infty \frac{e^{i(p_f - p_i)x}}{r - r_H} dx^4 \equiv if(u) \int_0^\infty \frac{e^{i(p_f - p_i)x}}{r - r_H} dx^4 \quad (16.5)$$

rescaling $r \rightarrow r' + r_H = r$ and $t \rightarrow t' + (r_H/c) = t$ to minimize the resonance energy in $p_f - p_i$. We then obtain:

$$S_{if} \equiv iZ \frac{1}{V} \sqrt{\frac{m^2}{E_f E_i}} \bar{u}(p_f, s_f) \gamma^0 u(p_i, s_i) e^{i r_H q} \int_0^\infty \frac{e^{i(p_f - p_i)x}}{r} dx^4 \equiv if(u) e^{i q r_H} \int_0^\infty \frac{e^{i(p_f - p_i)x}}{r} dx^4 \text{ For so:}$$

$$S_{if} \equiv if(u) \left[\int_0^\infty \frac{e^{i(p_f - p_i)(x' - r_H/2)}}{|x'|} dx'^4 - \int_0^\infty \frac{e^{i(p_f - p_i)(x' + r_H/2)}}{|x'|} dx'^4 \right] \approx if(u) e^{i(r_H/2)q} \left(\int_0^\infty \frac{e^{i(p_f - p_i)(x')}}{|x'|} dx'^4 - 2\pi(r_H)^3 \right)$$

(10.8.6)

Note that $\bar{u}(p_f, s_f) \gamma^0 u(p_i, s_i) = (1 - \beta \sin^2 \frac{\theta}{2})$ = Mott scattering term with the $e^{i(r_H/2)q}$ our

resonance term. The other left side coefficients and reciprocal $|x|$ part of S_{ij} comprise the well known Rutherford scattering term

$d\sigma/d\Omega = [(Z_1 Z_2 e^2)/(8\pi\epsilon_0 m v_o^2)]^2 \csc^4(\phi/2) = 1.6 \times 10^4 (\csc(\phi/2)/v_o)^4$. (Note that equation 10.8.6 applies to the $2P_{1/2}$ - $2P_{3/2}$ state electron-electron interaction (i.e., neutron) below). Here $p_f = p = q$. Note in equation 10.8.6 the factor $i e^{i k q} = i(\cos k q + i \sin k q)$. Here we find the rotational resonances at the $2P_{3/2}$ $r = r_H$ lobes associated with maximizing the imaginary part which is $i \cos k q$ to obtain absorption scattering (at $k q = \pi$), which here will then be the masses exchanged in inverse beta decay. Also a solution to the Dirac component is always a solution to equation 14.1 (but not vice versa) if we invoke an integer spin in this resonance term. Here also the p part uses the old De Broglie wave length to connect to the $p = h/\lambda$. In that regard recall that $h\nu/c = h/\lambda = p$ and for a DeBroglie wave fundamental harmonic resonance we have $\lambda_{rot} = 2\pi r$ for a stationary particle of **spin 1**= L (ambient E&M field source gives $L=1$ De Broglie).

Coulomb scattering of electrons, taking account spin-spin scattering

$$\frac{d\sigma}{d\Omega} = \frac{4Z^2 \alpha^2 m^2}{2|q|^4} Tr \gamma_o \frac{(\not{p}_i + m)}{2m} \gamma_o \frac{(p_i + m)}{2m}$$

$$\frac{d\sigma}{d\Omega} = \frac{Z^2 \alpha^2}{4p^2 \beta^2 \sin^4(\theta/2)} \left(1 - \beta^2 \sin^2 \frac{\theta}{2} \right)$$

Mott scattering, relativistic correction to Rutherford scattering

Ultrarelativistic electron scattering: Electron rest mass m neglected.

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4E^2} \left(\frac{(\cos^2(\theta/2)) - (q^2/(2M^2))(\sin^2(\theta/2))}{(\sin^4(\theta/2))(1 + (2E/M)\sin^2(\theta/2))} \right)$$

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4E^2} \left(\frac{(\cos^2(\theta/2)) - ((q^2/(2M^2))\sin^2(\theta/2))}{((\sin^4(\theta/2)))(1 + (2E/M)\sin^2(\theta/2))} \right) \quad (10.8.7)$$

$m/E \ll 1$, $m^2 \rightarrow 0$, $q^2 = (p_f - p_i)^2 = -4EE' \sin^2(\theta/2)$. Proton behaves like a heavy electron of mass M . $E \rightarrow 1/\sqrt{\kappa_{00}} = 1/\sqrt{(1 - \epsilon - \Delta\epsilon - r_H/r)}$.

For forward scattering $\theta \approx w/\beta^* \approx 0$ in $\sin^2(\theta/2) \ll 1$ in the below figure 6. So

$d\sigma/d\Omega = d(1/E^2)/dE^2 (1 + \text{tiny})/\text{tiny} = ((d(1/t)/dt)(1 + \text{tiny})/\text{tiny} = -(1/t^2)(1 + \text{tiny})/\text{tiny}$.

$t = E^2 = (\text{energy transfer})^2$.

Ultrarelativistic dependence of 10.8.7 new pde electron differential cross-section on $1/E^2$.

E =energy. Recall in this theory this should also be the energy dependence of ultrarelativistic proton-proton scattering since protons are made of electrons (in my work) and at very high

energies ($E \gg 150 \text{ GeV}$) the electron cloud binding energies in the proton don't matter anymore (that Paschen back binding energy starts becoming negligible at TeV energies): we have free electrons hitting free electrons once again.

For energy transfer t on left side graph (fig.6) $d\sigma/dt \propto d(1/E^2)/dE^2 = d(1/t)/dt = -1/t^2$, $t=E^2$. Energy transfer \sqrt{t} is proportional to $1/p$. But p^2 is proportional to area which is then proportional to $1/\Delta E^2 = 1/t$. So $\sigma = \text{Area} \propto 1/\Delta E^2 = 1/t$. $\Delta\sigma/\Delta\sqrt{s} = (1000\text{nb} - 10\text{nb})/1\text{GeV}$. But this is my equation 1 in figure 6 (also eq.10.8.7) for near forward elastic scattering. For $1\text{GeV} \approx 1\sqrt{s}$ then this $\Delta\sigma/(1)$ is a measure of $d\sigma/dt$ on the left side forward scattering elastic energy transfer graph since $1^2=1$.

But square root energy transfer \sqrt{t} in a scattering event for a beam at a specific energy (let's say at 13KeV) is also the abscissa of that big graph (on the left of the totem figure 6). So it is possible to get from total cross-section σ of electron scattering verse energy \sqrt{s} : σ/\sqrt{s} to $d\sigma/dt$ vs t where $t = (\text{energy transferred})^2$ at least at (literally) ONE (1GeV) energy transfer.

The fact that LHC totem measures elastic forward scattering thereby made it possible to test this theory (eq.2, 1.11, new pde) at 13TeV (and $\sqrt{s} \approx 2$), the very highest energy particles that mankind can produce. I could estimate from LHC data the asymptotically infinite beam energy transfer (curve) energy (red line) and compare it with my own σ/\sqrt{s} at $\sqrt{s} \approx 2$. From the graph of my equation 1 (10.8.7):

1000nb at $\sqrt{s} = 1.5\text{GeV}$

100nb at $\sqrt{s} = 2\text{GeV}$

10nb at $\sqrt{s} = 2.5\text{GeV}$

But that curve of eq.1 in figure 6 is for one eq.2 electron scattering off of one equation 2 (new pde) electron. Since there are 3 such electrons in each of the two protons you must multiply by 9 to get $d\sigma/dt$.

On my QED graph of my eq.1 had $\sigma \approx 100\text{nb}$ at about $2 \approx \sqrt{s}$. So multiply by 9 and get $\Delta\sigma/\Delta s \approx (1000\text{nb} - 10\text{nb})/((1.5 - 2.5)\sqrt{s}) = 10^{-3}\text{mb}/1\text{GeV}$. But $1^2=1$ and there are 3 electrons per proton so we multiply by 2: $9 \times 10^{-3}\text{mb}/1^2\text{GeV}^2 \approx 10^{-2}\text{mb}/1^2\text{GeV}^2$ which is sitting in approximately the same $t \approx 2$ spot on the left side $d\sigma/dt$ vs t graph of Fig.6.

So we proved from the data that a ultrarelativistic proton-proton scattering event ($\sim 13\text{TeV}$) is equivalent to 6 free electrons scattering off each other with the electrons obeying (2AI) equation 9, the new pde. Thus the hadron theory that should be used is $2+2+2$ at $r=r_H$, not quark theory.

Note also the cusp is at the proton reduced mass here. It is where ($\sim .5\text{GeV}$) binding energy must be added to break the electrons apart in the head on collision which takes away from the elastic scattering energy transfer. So we must apply this theory at much higher energies (eg., $\sqrt{s} \approx 2$).

Quark theory (QCD) implies some kind of exponential dependence which is not seen in this scattering data.

Hard Shell Scattering Peak Of $d\sigma/ds$ Implies Protons Made of Electrons, not Quarks

The electron radius at $2.8 \times 10^{-15} (m_e/m_t) = 8.1 \times 10^{-19}\text{m}$ provides the hard shell cross-section limit.

For colliding beams we have an additional factor of 2 here. $2(m_\tau + m_\mu)/m_e = 6.91347\text{TeV}$.

There 3 are electrons in the proton so the proton energy is $3 \times 6.913\text{TeV} = 20.74\text{TeV} \approx 21\text{TeV}$. So the $d\sigma/ds$ should level off at proton energy 21TeV . This is in analogy with the $Q = \sigma/\pi r^2 \propto d\sigma/ds$ ($1/s \propto \lambda$) $r = \lambda$ peak of Mie scattering theory.





Fig6
Analogy to Mie scattering



Extinction efficiency (Q_e) as a function of diffraction parameter $x (= 2\pi r/\lambda)$.

Analogy of Mie scattering with $Q = \sigma/\pi r^2$. Here the lepton hard shell is at $r = 2 \times 10^{-19} \text{ m}$. Note analogy of leveling off of $d\sigma/ds$ $\lambda = r$ (i.e., $x=6$) as at $x=21 \text{ TeV}$ for LHC.

Totem

LHC totem forward scattering gives elastic scattering cross-sections for high energies and so small scattering angles. We choose $\theta = 1^\circ$.

$$\left(\frac{d\sigma}{d\Omega}\right)_B = (3) \frac{\alpha^2}{8E^2} \left[\frac{1+\cos^4(\theta/2)}{\sin^4(\theta/2)} - \frac{2\cos^4(\theta/2)}{\sin^2(\theta/2)} + \frac{(1+\cos^2\theta)}{2} \right] =$$

$$1.43 \times 10^{-43} \left(\frac{1 - .99985}{5.8 \times 10^{-9}} - \frac{2(.99985)}{7.6 \times 10^{-5}} + \frac{1 + .9997}{2} \right)$$

$$= 1.43 \times 10^{-43} (2.626 \times 10^4 - 2.631 \times 10^4 + .99985) = (1.43 \times 10^{-43}) 52.788$$

$$= 7.55 \times 10^{-42} = \mathbf{7.55 \times 10^{-14} \text{ barns/steradian.}}$$

This result might be found in the totem data archives.

LHC totem forward scattering gives Coulomb scattering cross-sections for high energies for larger (but still small) scattering angles. We choose $\theta=3^\circ$. So

$$\frac{d\sigma}{d\Omega} = (3) \frac{Z^2 \alpha^2}{4p^2 \beta^2 \sin^4(\theta/2)} \left(1 - \beta^2 \sin^2 \frac{\theta}{2} \right) =$$

$$= (3) \frac{(1) 5.33 \times 10^{-5}}{4(1.04 \times 10^{37} 4.7 \times 10^{-7})} (1 - 1^2 6.85 \times 10^{-4}) = 8.18 \times 10^{-38} m^2 =$$

$$= \mathbf{8.17 \times 10^{-10} \text{ barns/steradian.}}$$
 This result might be found in totem data archives data.

Meson Multiplets

"tetra quarks" are merely two mesons bound together! They can bind together more deeply if the components of the mesons themselves are bound individually to the components of the other meson giving more mass, section 8.11.

In this theory (DavidMaker.com, Ch.9-10) this is called singlet and doublet states with one bound with more binding energy than the other for those heavy upper 2P Paschen Back states. So these look like heavy and light tetraquark states but they are not, they are merely two types of meson binding states. You could predict the energies from the Paschen Back effect associated with those large plate fields, section 8.11.

References

Pugh, Pugh, *Principles of Electricity and Magnetism*, 2nd Ed. Addison Wesley, pp.270
 Bjorken and Drell, *Relativistic Quantum Mechanics*, pp.60
 Sokolnikoff, *Tensor Analysis*, pp.304

11.1. W Compton Wavelength Region

Recall in appendix A m_e Source Term at $r=r_H$ Inside Angle C.

Analogously from **2AC** we get with the eq.3 doublet $\varepsilon \pm \varepsilon$ the Proca equ (3) *neutrino and electron* $\Delta \varepsilon$ at $r=r_H$. As in sect.6.13 in κ_{00} we normalize out the muon ε . So we are left with the electron $\Delta \varepsilon$ in $\kappa_{00} = 1 - [\Delta \varepsilon / (1 \pm 2\varepsilon)] + [r_H (1 + ((\varepsilon \pm \varepsilon)/2)) / r]$ from the two above rightmost (Proca) diagrams. So Source = $E_{ZW} = \frac{1}{\sqrt{\kappa_{00}}} = \frac{1}{\sqrt{1 - \frac{\Delta \varepsilon}{1 \pm 2\varepsilon} - \frac{r_H \varepsilon (1 + (\varepsilon \mp \varepsilon)/2)}{r}}} \approx \frac{1}{(1 \pm \varepsilon) \sqrt{\Delta \varepsilon}}$, at $r = r_{H_e} +$ is for Z

and $-$ is for W. So W (right fig.) is a single electron $\Delta \varepsilon + v$ perturbation at $r=r_H=\lambda$ (since m_e ultra relativistic): So $H=H_0+m_e c^2$ inside V_w . $E_w=2hf=2hc/\lambda$, $(4\pi/3)\lambda^3=V_w$. For the two leptons $\frac{1}{V^{1/2}} = \psi_e = \psi_3, \frac{1}{V^{1/2}} = \psi_\nu = \psi_4$. Fermi $4pt = 2G \iiint_0^{r_w} \psi_1 \psi_2 \psi_3 \psi_4 dV = 2G \iiint_0^{r_w} \psi_1 \psi_2 \frac{1}{V^{1/2}} \frac{1}{V^{1/2}} V = 2 \iiint_0^{r_w} \psi_1 \psi_2 G \equiv \iiint_0^{V_w} \psi_1 \psi_2 (2m_e c^2) dV_w = \iiint_0^{V_w} \psi_1 (2m_e c^2) \psi_2 dV_w$. (A3)

What is Fermi G? $2m_e c^2 (V_w) = .9 \times 10^{-4} \text{ Mev} \cdot F^3 = G_F$ **the strength of the weak interaction.**

Derivation of the Standard Model But With No Free Parameters

Since we have now derived M_W , M_Z , and their associated Proca equations, m_μ, m_τ, m_e , etc., Dirac equation figure 2 part.1), G_F , ke^2 , Bu, Maxwell's equations, etc. we can now set up a Lagrangian density that implies all these results. In this Formulation $M_Z = M_W / \cos\theta_W$, so you find the Weinberg angle θ_W , $g \sin\theta_W = e$, $g' \cos\theta_W = e$; solve for g and g' , etc., We will have thereby derived the standard model from first principles (i.e., postulate 1) and so it no longer contains free parameters!

Thus we have the interaction ε operating in W radius using the doublet of Ch.3.

In general then we have obtained an ortho triplet state here since we are merely writing the Clebsch Gordon coefficients for this addition of two spin $1/2$ angular momentums:

$|1/2, 1/2, 0, -1\rangle, \dots |1/2, 1/2, 0, 0\rangle, \dots |1/2, 1/2, 0, 1\rangle, \dots$ or $+W, Z, -W$.

Anyway, this small S matrix involves the neutrino and so can allow spin $1/2$ neutrino emission jumps instead of just the usual E&M spin 1 jumps. 100km/sec metric quantization translates to a neutrino rest mass of .165eV.

11.2 Excited Z States

Put m_e in Equation 6.4.1

The beautiful thing to be noted here is that for the doublet resonance with the $2P_{3/2}$ lobe at $r=r_H$ that minimizes energy you get the spin 1 W and Z and the value of the Fermi G ! We have also shown that this doublet interaction corresponds to the exchange of massive spin 1 particles (recall spin $1/2$ s forbidden by that $j-1/2$ factor).

11.3 Probability for $2P_{3/2}$ Giving One Decay 1S Product at $r \approx r_H$ In W Region

In equation 4.12 we note that invariance over 2π rotations using $(1+2\varepsilon)d^2\theta$ does not occur anymore thus seemingly violating the conservation of angular momentum. To preserve the conservation of angular momentum the additional angle ε must then include its own angular momentum conservation law here meaning intrinsic spin $1/2$ angular momentum in the S state case and/or isospin conservation in the $2P_{3/2}$ case at $r=r_H$. In any event we must also integrate to $C=\varepsilon$. Here we do the E&M component decay given by equation 3.2.

Plug in $S_{1/2} \propto e^{i\phi/2}$, $1/2(1-\gamma^5)\psi = \chi$ into the 4pt. interaction integral. In that regard note that the expectation value of γ^5 is proportional to $v \propto$ Heisenberg equation of motion derivative of $2P_{3/2} \propto e^{i(3/2)\phi}$. We integrate $\langle \text{lepton} | \text{baryon} \rangle$ over this W exchange region where we note $(\sim 1/100)F$ for 90Gev particle, so $dV = ((1/100)F)^3 = \text{Vol}W$. Also $ck_0 = \varepsilon = 106\text{Mev}$ from section 2.1.

From Ch.3 on the vacuum constituents e and ν we note that $\int d\tau = \text{Vol}$, χ is defined as the vacuum eigenfunction. Vacuum expectation sect.B2: $\Sigma \langle |v_{aM}\rangle | \varepsilon | \langle v_{aM+1} \rangle = \langle |$

$\int \psi^* \nu \varepsilon \chi_e dV \rangle = \langle | \text{Pot} \rangle = \varepsilon \text{Volume of } W$. Recall also that appendix A implies that the W and the Z are composites. This application of eq.2 for example applies to the $2P_{1/2} - 2P_{3/2}$ electron-electron scattering state inside the neutron $\langle \text{Proton} 2P_{3/2} | \text{Pot} | \text{Neutron} 2P_{1/2} - 2P_{3/2} \rangle$. Plug in $S_{1/2} \propto e^{i\phi/2}$, $1/2(1-\gamma^5)\psi = \chi$ Also we can get a weak, strangeness changing (second term below), decay from a $2P_{1/2} - 2P_{3/2} \rightarrow m_p$ to the S state branch equation. eq.2 expectation values in the 4pt..

$= \Sigma \langle \text{lepton} | \text{vac} \rangle | \varepsilon | \langle \text{vac} | \text{baryon} \rangle = \text{Fermi interaction integral} = \int \psi_1^* \psi_2 \psi_3 k_0 c \chi dV =$

$\int \psi_1^* (\varepsilon \text{Vol}W) \chi dV = \int \psi_1^* (\varepsilon \text{Vol}W) \chi dV$. Also $dV = dA d\phi = K d\phi$.

So the square root of the probability of being in the final state is equal to the Fermi integral =

$\int \psi_1^* (\text{Pot}) \chi dV = \int \psi_1^* \psi_2 \psi_3 \varepsilon \Delta V_W \chi dV =$

$$\begin{aligned}
&= \int K \langle e^{i\phi/2} | (\varepsilon \Delta V_W) | 1 - (\gamma^5 i e^{i(3/2)\phi}) \rangle dt dV = \int K \left[\varepsilon \left(\frac{F}{100} \right)^3 \right] \langle e^{i\phi/2} - \gamma^5 i e^{i4/2\phi} \rangle d\phi \\
&\varepsilon Vol W = \int K \left[\varepsilon \left(\frac{F}{100} \right)^3 \right] \langle e^{i\phi/2} - \gamma^5 i e^{i4/2\phi} \rangle d\phi \\
&= KG_F \int \langle e^{i\phi/2} - \gamma^5 i e^{i4/2\phi} \rangle d\phi = KG_F \left(\frac{2e^{i\phi}}{i} \Big|_0^{2\pi+C} - \frac{2\gamma^5 e^{i4\phi}}{i4} \Big|_0^{2\pi+C} \right) \quad (10.8.7)
\end{aligned}$$

with $\langle \text{initial} | c | \text{final} \rangle^2 \approx$ transition probability as in associated production with the separate 2P proton ground state transition being the identity ($\Delta S=0$). Factoring out the 2 and then normalizing 1 to .97 simultaneously normalizes the 1/4 to .24 in section 3.2. With this normalization we can set $\cos\theta_c=.97$ and $\sin\theta_c=.24$. Thus we can identify θ_c with the *Cabbibo angle and we have derived its value*. We can then write in the weak current sources for hadron decay the **VA structure**: $|\cos\theta_c - \gamma^5 \sin\theta_c|$. Thus with the above Cabbibo angle and this CP violation and higher order $(r_H/r)^n$ terms in section 3 we have all the components of the CKM matrix. Note we have also derived the weak interaction constant G_F here.

Given the role ε plays here in decay we find the expectation value of energy ε within the S matrix scattering region in chapter 10.

Recall from section 1.2 the possible mixing of real and imaginary terms in that energy coming out of that first order Taylor expansion. There we found the $1+x$ and $1-x$ solutions cancel and we could ignore the $1+1=2$ term as it is still a flat metric.

Also there are still extra terms provided by the 'small' higher order r^2 terms in that Taylor expansion so that "higher and lower" than the speed of light mixed condition still can exist (for $\Delta G \neq 0$. See end of section 4.6 and 10.8.6). In that regard note for the next higher order Taylor term at largest curvature $d^2(1/k_\pi)/dr^2$ is large negative and r^2 is positive implying a net negative term and therefore a neutral charge (see case 2, of section 19.6)! In that case the perturbative squared r term appears to overwhelm the rest since the lower order terms then cancel. Note from the above we put these neutral conditions also into that decay since net charge is zero in the Cabbibo angle derivation. This then appears to be the beta decay condition *where the neutrino (higher than c) and the electron, (lower than c), decay from this neutral particle condition* (bottom of section 10.8). The beginning $2P_{3/2}$ ground state still exists however in the respective Cabbibo angle calculation. Thus those real and imaginary terms coming out of that Taylor expansion provide the explanation for beta decay.




Fig.7

m_e Source Term Inside Angle

.See section 0.2 and B2 So W is a single electron $\Delta\epsilon$, v perturbation at $r=r_H$: $H=H_0+2m_e c^2$ inside V_w . $E_w=2hf=2hc/\lambda$, $(4\pi/3)\lambda^3=V_w$. For the two leptons $\frac{1}{v-1/2} = \psi_e = \psi_3, \frac{1}{v-1/2} = \psi_v = \psi_4$.

$$\text{Fermi } 4\text{pt} = G \iiint_0^{V_W} \psi_1 \psi_2 \psi_3 \psi_4 dV = G \iiint_0^{V_W} \psi_1 \psi_2 \frac{1}{V^{1/2}} \frac{1}{V^{1/2}} V = \iiint_0^{V_W} \psi_1 \psi_2 G \equiv \iiint_0^{V_W} \psi_1 \psi_2 (2m_e c^2) dV_W. \quad (\text{A3})$$

What is Fermi G? $2m_e c^2 (V_W)/F^3 = .9 \times 10^{-4} \text{MeV} \cdot F^3 = G_F$ **the strength of the weak interaction.**

Next we plug the respective ψ s into ψ_1, ψ_2 in sect.B2. In that regard the expectation value of γ^5 is speed and varies with $e^{i3\phi/2}$ in the trifold. The spin $1/2$ decay proton $S_{1/2} \propto e^{i\phi/2} \equiv \psi_1$, the original $2P_{1/2}$ particle is chiral $\chi = \psi_2 \equiv 1/2(1 - \gamma^5)\psi = 1/2(1 - \gamma^5 e^{i3\phi/2})\psi$. Initial $2P_{1/2}$ electron ψ is constant. Plug these terms into equation B2 $= \iiint_0^{V_W} \psi_1 \psi_2 (2m_e c^2) dV_W = \iint \psi_{S1/2}^* (2m_e c^2 V_W) \chi dV =$

$$K \int \langle e^{\frac{i\phi}{2}} [\Delta \varepsilon V_W] \left(1 - \gamma^5 e^{i\phi \frac{3}{2}}\right) \psi \rangle d\phi = K G_F \int \langle e^{i\phi/2} - \gamma^5 i e^{i4/2\phi} \rangle d\phi = K G_F \left(\frac{2e^{i\phi}}{i} \Big|_0^{2\pi+C} - \frac{2\gamma^5 e^{i4\phi}}{i4} \Big|_0^{2\pi+C} \right)$$

with $VA \langle \text{initial} | c | \text{final} \rangle^2 \approx$ transition probability as in associated production. Factoring out the 2 and then normalizing 1 to .97 simultaneously normalizes the 1/4 to .24 in Ch.3. With this normalization we can set $\cos\theta_c = .97$ and $\sin\theta_c = .24$. Thus we can identify θ_c with the **Cabbibo angle and we have derived its value.**

$$\Gamma = [G^2/(4\pi N)] m_i^\alpha P_i^{|\mathbf{m}|} |\mathbf{x}|^2, \quad \mathbf{x} = \Sigma \mathbf{m}_j / m_i \quad \text{Eg. } 1/\tau_\mu = [G^2 m_\mu^5 / (192\pi^3)] (1 - m_e^2/m_\mu^2)^6.$$

r<rh Application: Rotational Selfsimilarity With pde Spin: CP violation

12.1 Fractal selfsimilar spin

The fractal selfsimilarity with the spin in the (new) Dirac equation 2 implies a selfsimilar cosmological ambient metric (Kerr metric) rotation as well as in section 4.1. Thus there will be $2ds, ds_\phi$ rotation metric cross terms with the dt (without the square) implying time T reversal nonconservation and therefore CP *non*conservation since CPT is always conserved. We thereby derive CP nonconservation from first principles: **CP nonconservation is a direct consequence of the fractalness.** This adds another matrix element of magnitude $\sim 1/3800$ (sect.6.3) for Kaon decays thus adding off diagonal elements to the CKM matrix.

Or for Kerr rotator use

$$ds^2 = \rho^2 \left(\frac{dr^2}{\Delta} + d\theta^2 \right) + (r^2 + a^2) \sin^2 \theta d\phi^2 - c^2 dt^2 + \frac{2mr}{\rho^2} (a \sin^2 \theta d\theta - c dt)^2, \quad (13.1)$$

$$\rho^2(r, \theta) \equiv r^2 + a^2 \cos^2 \theta; \quad \Delta(r) \equiv r^2 - 2mr + a^2, \text{ or } ds^2 = dr^2 + dt^2 + 2dt dr + ..$$

In a polarized state ($\theta = 0^\circ, 180^\circ$) in 25.3, 25.25 the off diagonal elements are proportional to $\phi = (\phi + c)e^{-C}$. Thus if the charge e is conjugated (C , e changes sign), if dr changes sign (P , parity changes sign) and dt is reversed (t reversal) then the ds quantity on the left side of equation 1.6 is invariant. But if dr (P) changes sign by itself, or even e and P together (CP) change sign then ds is not invariant and this explains, in terms of our fractal picture, why CP and P are not conserved generally. P becomes maximally nonconserved in weak decays as we saw in above. The degree to which this nonconservation occurs depends on the “ a ” (in eq.3.2.1) transfer $\langle \text{final} | a | \text{initial} \rangle$ (equation 3.2) which itself depends on the how much momentum and energy is transferred from the S_{M+2} to the S_{M+1} fractal scales as we saw in this section. Recall chapter 5 alternative derivation of that new (dirac) equation pde (eq.2) **linearization of the Klein Gordon equation** ($c=1, \hbar=1, m=1$, eq.2):

$$\left(-\alpha_1 i \frac{\partial}{\partial x_1} - \alpha_2 i \frac{\partial}{\partial x_2} - \alpha_3 i \frac{\partial}{\partial x_3} + \beta \right) \left(-\alpha_1 i \frac{\partial}{\partial x_1} - \alpha_2 i \frac{\partial}{\partial x_2} - \alpha_3 i \frac{\partial}{\partial x_3} + \beta \right) = \quad (13.2)$$

$$-\alpha_1^2 \frac{\partial^2}{\partial x_1^2} - \alpha_2^2 \frac{\partial^2}{\partial x_2^2} - \alpha_3^2 \frac{\partial^2}{\partial x_3^2} + \beta^2 + 2 \sum_{\mu=1}^3 \sum_{\nu=1}^3 \alpha_\mu \alpha_\nu \frac{\partial}{\partial x_\mu} \frac{\partial}{\partial x_\nu} + 2 \sum_{\mu=1}^3 \beta i \alpha_\mu = -\frac{\partial^2}{\partial t^2}$$
 This equals $= c^2 p_1^2 + c^2 p_2^2 + c^2 p_3^2 + m^2 c^4 = E^2$ if the off diagonal elements zero which is the condition used in the standard Dirac equation derivation of the α s and β . Note that the off diagonal elements $2 \sum_{\mu=1}^3 \sum_{\nu=1}^3 \alpha_\mu \alpha_\nu \frac{\partial}{\partial x_\mu} \frac{\partial}{\partial x_\nu} + 2 \sum_{\mu=1}^3 \beta i \alpha_\mu$ are equivalent to the off diagonal elements in equation 5.1 (and are corrections to 5.2 in fact) so *are not* zero for parity and CP NONconservation in this context (in a rotating universe). So in the context of the Dirac equation the CP violation term $e_s(dr)dt \rightarrow (dr/ds)(dt/ds) \rightarrow pE\chi$ (after division by ds^2). Thus CP violation goes up as the square (pE) of the energy (so should be larger in bottom factories). The section 13.2 below Cabbibo angle calculation (not rotation related however) is an example of how this method can give the values of the other terms in the CKM matrix. They arise from calculation of $\langle Z \rangle$ between higher order m harmonics.

This section is important in that we see that CP violation is explainable and calculable in terms of perturbative effects on the ambient metric (and therefore the Dirac equation) of a rotating universe with nearly complete inertial frame dragging (eq. A6 in the E&M form), CP violation doesn't need yet more postulates as is the case with the GSW model. In fact the whole CKM matrix is explainable here as a consequence of this perturbation.

Note the orientation relative to the cosmological spin axis is important in CP violation. Integration of the data over a 3 month time (at time intervals separated by a sidereal day) is going to yield different CP violation parameters than if integration is done over a year.

Miscellaneous

12.2 GIM Derivation

Recall in the GIM (Glashow, Iliopoulos, Maiani) hypothesis that u,d were a pair of left handed Fermion states as in V-A . $d' = d \cos \theta_c + s \sin \theta_c$, $s' = -d \sin \theta_c + s \cos \theta_c$ where θ_c is the Cabbibo angle. Thus u,d are paired, s,c are paired, b,t are paired and we have the V-A transitions.

Here we identify the new pde 2P state for $r=r_H$ has P_x , P_y and P_z states which split in energy due to that Paschen Back effect given those ultrarelativistic plates, into paired spin up and spin down states (P_x, P_x') , (P_y, P_y') , (P_z, P_z') analogous to the GIM (u,d),(s,c),(b,t). Here the spin orbit interaction (LS) coupling energy term is much stronger than the SS coupling term. So we have pairs of states J, M, M' with P_x and P_y being orthogonal, except for those weak interaction V-A terms. The ds^2 to ds' transition is through the V-A term. Recall equation the 16.7 $|\int \chi_f^* G \chi_o dV|^2 =$ =transitionprobability of a ds^2 to a ds . of eq.B2 ($\chi = .5(1-\gamma^5)\psi$ with ψ in ds^2 , χ in ds) for V-A Cabibbo angle transitions (transitions inside P_x separately from P_y and separately from P_z) where $|1| - |\gamma^5|$ replaces the $\cos \theta_c + \sin \theta_c$. So in analogy for transitions between P_x and P_y $P_x' \cos \theta_c + P_y \sin \theta_c = P_x'$, $P_y' = -P_x' \sin \theta_c + P_y \cos \theta_c$. This is a first principles understanding of GIM thereby allowing us to derive the electroweak cross-sections (WS).

Recall that $dz = -1, 0$ solution to eq.2 for $C=0$ implies $dr < 0$ at least for small C. (low noise). because -1 is on the real r axis.

12.3 Normed Division Algebra, Octonians, E8XE8 and SU(3)XSU(2)XU(1) Basis Change

Note from the above that the new pde fractal theory generated the electron 2AI with mass, the near zero mass left handed neutrino 1.12. Recall also from above the $\kappa_{00}=\sqrt{1-\epsilon-r_H/r}$. The W was generated from a nonzero ambient metric ϵ in that S matrix derivation part of the metric coefficient $\kappa_{\mu\nu}$. Interestingly that Normed Division Algebra (NDAR) on the real numbers (as in: $\|Z_1\|*\|Z_2\|=\|Z\|$) from equation 1 implies that octonians (and thereby the *largest normal Lie group* E8XE8) are also allowed. Recall we have that SU(2) Lie group rotation for the 0° extrema imbedded in a E8XE8 rotation since one of its subgroups being SU(3)XSU(2)XU(1). This is the only subgroup we can use because it is the one that only contains that SU(2).

Eigenstates

Recall the $m_\tau=1$ was separable at 45° from the rest (of the eq.1.15 diagonal states) since ds is constant there for small rotations. So ds_τ can be normalized.

The B field rotations are here reciprocals of the rotations in the Mandelbulbs since $\kappa_{00}=1-r_H/r$ and so $r_H/r \rightarrow \xi dr$ for B field motion given smaller radius r means high energy ξdr . So the *ortho* state is the smaller ϵ Mandelbulb eigenstate and the *para* state is the larger τ limaçon Mandelbulb eigenstate.

Meta Theory Of Couplings In SM

From the 1.15 diagonal on the Mandelbrot set: $m=m_L=1+\epsilon+i\Delta\epsilon=m_\tau+m_\mu+m_e$.

$$mv^2/r_H=qvB \quad (1)$$

$$\pi r_H^2 B = \Phi_0 \quad (2)$$

$v=c$. Solve Eq.1 and eq.2 for q :

$$q=mc\pi r_H/\Phi_0. \quad (3)$$

The effect of the *E field lines coming together* by Fitzgerald contraction γ imply a force increase that can be realized by invoking an *effective* charge increase $e \rightarrow q'$. or in $V/dr' = \text{Electric field}$ with $r=r_H$ in $dr'^2 = \kappa_{rr} dr^2 = [1/(1-r_H/r)] dr^2$ (The charge e itself really does not change.).

For $m=m_\mu = \epsilon$ 2P_{3/2} state (So ultrarelativistic so E field line contraction.). **From equation 3**

$q' = e$, $H \rightarrow e^2$ E&M for the Nth fractal scale, **Gravity** for the N+1th fractal scale.

For $m=m_\tau + m_e$ as meson. 2P_{3/2} state (so ultrarelativistic). **From equation 3:**

$q' = 46e$, $H \rightarrow (46e)^2$ Strong Force.

For $m=m_e = i\Delta\epsilon + v$, v small, 2P_{1/2}, $dr'^2 = \kappa_{rr} dr^2 = 1/(1-r_H/r) dr^2$. So $V/dr' = E$ small. **From equation 3:**

$q' = ie/200$, $H \rightarrow iq'V$ so $\psi(t) = e^{iq'Vt}\psi(t_0) = e^{-q'Vt}\psi(t_0)$.

exponential decay with a force $q'^2 = 40000X$ *smaller* than the E&M. **Weak interaction.**

dr' large allowing large uncertainty principle dr' for small nonrelativistic mass m_e in $(dr' \bullet m_e c) \geq \hbar/2$. This occurs for small externally observed dr and $m_e c$ in the 2P_{1/2} state and 1S_{1/2} state at $r=r_H$. But these are decay states (PartII Sect.7.3). Given these strength and decay parameters we can alternatively integrate over the r_c volume our W and Z particles to get the Fermi G 4pt coupling of weak interaction theory in the SM. W is then a virtual intermediary here. **So we just derived all 4 forces from that diagonal on the Mandelbrot set.**

Calculations: So for the Kerr mass ortho state (2nd Mandelbulb) $(a/r)^2 = \epsilon + \Delta\epsilon$ (thus added to 1)

at $r=r_H$, for (N+1): $m_e v^2/r = qvB$ so $m_e v/(qB) = (1-2\epsilon)m_\mu c/qB = r$. Thus $(1-2\epsilon)m_\mu c/q = rB =$

$$(1-2\epsilon)(1.883 \times 10^{-28})(299792458)/(1.6022 \times 10^{-19}) = .3525(1-2\epsilon)$$

$$\Phi_0/(rB) = \Phi_0/[0.3525(1-2\epsilon)] = B\pi r^2/(rB) = \pi r.$$

$$r_H = 1.359 \times 10^{-15}(2)/[(.3526)\pi(1-2\epsilon)] = 2.805 \times 10^{-15} m = e^2/m_e c^2 \text{ for 2P states (eq.7.1).}$$

Compare and contrast with the mainstream Toy Model

12.4 Derivation of Mainstream Toy Model Composite System $|1\rangle|2\rangle|3\rangle$

Interpretation From Newpde 3e State

To explain the above composite 3e stability result sect.2 implies $r=r_{HP}$ in $dt'=0$ in $dt'^2=\kappa_{00}dt^2=(1-r_{HP}/r)dt^2$ (in the new pde, nucleon radius) and so $dt'=0$ so that clocks stop. So we have complete **proton stability** in the new pde given eq. 11 $2P_{3/2}$ at $r=r_{HP}$ fills first (see review section just above).



$2P_{3/2}$ is trifolium shaped $\psi^*\psi$ so the electron spends 1/3 time in each lobe (**fractional charge**), lobes can't leave (**asymptotic freedom**), P wave scattering (**jets**), 6 P states (**6flavors uds c b t**) **explaining the major properties of quarks** and so explaining the strong interaction. Note that $ds_1^2 \neq 0$ with $dt \approx 0$ so $\sqrt{2}ds^2 = dr^2 + dt^2$ implies, after eq.3.6 operator formalism derivatives are put in (figure 4 2nd diagram from right), the Klein Gordon equation spin 0 mesons, the carriers of the strong force. Also our Mandelbulb analysis implies that the proton mass is 937Mev (appendix C) implying ultrarelativistic internal electron motion is needed to get this mass. Given this ultrarelativistic electron (needed to obtain the much larger proton mass hard shell result P. Alberto, R. Lisboa, M. Malheiro and A. S. de Castro, Phys.Rev. 58 (1998) R628) at r_H the field lines must be Fitzgerald contracted to a flat "plate" and thus be high density, field lines thus **explaining the strength of the 'strong' force**.

$2P_{3/2}$ fills first in Aufbau principle for ultrarelativistic hard shell (Alfredo 1998).

Electron in limaçon lobe added to trifolium lobe to give bound state:

$-e + (2/3)e = -(1/3)e$ d. Add other two lobes $(2/3)e$ u + $(2/3)e$ u = uud =

Fill inrest of P states same way.



6 P orbital slots at $r=r_H$ Fill states as nondegenerate energy (level) goes up \longrightarrow

Possible SHM interaction between these lobes gives excited states.

LS coupling Lande' g-factor structure gives minimal LS energy for smallest L

So net spin 1/2 states preferred.

Note here we have three ultrarelativistic $2P_{3/2}$ electrons (2 positrons and one electron) needed to create the proton (which as we noted above is much heavier) at $r=r_H$. Given a central (negative) electron the two outer positron $2P_{3/2}$ state plates (at 120°) only intersect at the center and so don't see each other at all and so don't repel each other, **explaining why the proton is still a bound state even though the two positive positrons are both inside r_H** (along with that electron).

Possible (but low probability) positron-electron annihilation inside r_H also implies, given the momentum transfer to the *third particle* with strong field plate and large mass (quadruply differential cross-section), that the resulting gamma ray will be short lived and pair creation will occur almost immediately within r_H , since $\sigma=1/20$ barn $\approx \pi r_H^2$ the cross-sectional equatorial area of the proton, guaranteeing pair creation occurs) replacing the previous pair immediately.

12.4 Single Electron Probability Trifolium Statistics Inside r_H

A single electron in the trifold implies that on average each of the 3 trifold lobes has $(1/3)e$ charge (hence the origin of hyperon fractional charge of the lobes). This allows for a toy model in which we give these $\psi^*\psi$ $2P_{3/2}$ at $r=r_H$ lobes (not particles) names (quarks, the toys.).

Clebsch Gordon Coefficients For Newpde state: $2P_{3/2}$ at $r=r_H$

It is well known that (and also implied by the new pde eq.11 with C_M) for the composite system of two electrons $|1\rangle|2\rangle$ you get, from the analysis of the invariance of the resulting Casimir operator J^2 , the resulting state $|J_A, J_B, J, M\rangle$ with combined operator $J_A + J_B = J$. This is our para state t and 3 ortho states below. For the third spin $1/2$ particle of far lower energy (the central electron, object B) we have $|1\rangle|2\rangle|3\rangle$ and so the Clebsch Gordon coefficients imply the decomposition $(2\otimes 2)\otimes 2 = (3\otimes 2)\oplus(1\otimes 2) = 4\oplus 2\oplus 2$ so that **three spin 1/2 particles** group together into **four spin 3/2** and only two spin $1/2$ s, **6 states** altogether, (the splitting u,d,s,c,b,t). Note then the **majority $2P_{3/2}$** (trifold core) states. Recall also that the $2P_{3/2}$ solution to new pde at $r=r_H$ gives a trifold shape, and $2P_{3/2}$ fills first. This fills in the broken degeneracy ortho states at the end of part II. The states close to the proton mass are filled in by the Frobenius solution below.

Eq.2 Single Electron Probability (trifold statistics):

Consistent with the toy model and also the electron or positron moving between lobes, this time using integer charge distributed over all three $2P_{3/2}$ lobes at $r=r_H$, just randomly put the lobe charges (lobe,lobe,lobe) on top of one another Monte Carlo style to determine the probability of a given charge in each lobe. For two positrons $[(+1/3,+1/3,+1/3)+(1/3,+1/3,+1/3)]$ and one electron $(-1/3,-1/3,-1/3)$ ($2P_{3/2}$) the probability of seeing a $+(2/3)e$ lobe is twice that of seeing a $-(1/3)e$ lobe so $((2/3,2/3,-1/3$ or uud proton) eg.,proton, C and b are the $1/2$ state components of $(2\otimes 2)\otimes 2$. For $(2P_{3/2})$ two positrons $[(+1/3,+1/3,+1/3)+(1/3,+1/3,+1/3)]$, an electron $(-1/3,-1/3,-1/3)$ and an outlier electron ($2P_{1/2}$) $(-1/3,-1/3,-1/3)$ the probability of seeing a $-(1/3)e$ lobe is twice as high as a $+(2/3)e$ lobe so $(-1/3,-1/3,2/3)$ or ddu Neutron).

It is well known that (and also implied by the new pde) for the composite system of two electrons $|1\rangle|2\rangle$ you get, from the analysis of the invariance of the resulting Casimir operator J^2 , the resulting state $|J_A, J_B, J, M\rangle$ with combined operator $J_A + J_B = J$. Using the resulting Clebsch Gordon coefficients we find the decomposition $2\otimes 2 = 3\oplus 1$, $m=1,0,-1$ ortho triplet state and singlet para state, which indeed are well known.(eg., Zeeman or Paschen Back line splitting, Ch.8.).

But for a third spin $1/2$ particle we have $|1\rangle|2\rangle|3\rangle$ and so the Clebsch Gordon coefficients imply the decomposition $(2\otimes 2)\otimes 2 = (3\otimes 2)\oplus(1\otimes 2) = 4\oplus 2\oplus 2$ so that **three spin 1/2 particles** group together into **four spin 3/2** and only two spin $1/2$ s, **6 states** altogether. Note then the majority $2P_{3/2}$ (trifold core) states.

We could now quit and use the mainstream quark (our $2P$ lobes) applications but that theory is inadequate (eg., neutron-proton binding energy, sect.10.7) and instead will proceed to directly solve equation 2 using the Frobenius series method.

Arfken, *Mathematical Methods of Physics*, 3rd ed. Page 454

Engel, Harold, *Introduction To Nuclear Physics*, 1966, Addison Wesley, page. 45

