

It's Broken, fix it

David Maker

Key words, Mandelbrot set, Dirac equation, Metric

Abstract In that regard Dirac in 1928 made his equation(1) flat space(2). But space is not in general flat, there are forces.

So over the past 100 years people have had to try to make up for that mistake by adding ad hoc convoluted gauge force after gauge force until fundamental theoretical physics became a mass of confusion, a train wreck, a junk pile. So all they can do for ever and ever is to rearrange that junk pile with *zero actual progress* in the most fundamental theoretical physics* ,.. forever. We died.

By the way note that Newpde(3) $\gamma^\mu \sqrt{\kappa_{\mu\mu}} \partial\psi/\partial x_\mu = (\omega/c)\psi$ is NOT flat space (4) so it cures this problem (5).

References

(1) $\gamma^\mu \partial\psi/\partial x_\mu = (\omega/c)\psi$

(2) Spherical symmetry: $(\gamma^x \sqrt{\kappa_{xx}}dx + \gamma^y \sqrt{\kappa_{yy}}dy + \gamma^z \sqrt{\kappa_{zz}}dz + \gamma^t \sqrt{\kappa_{tt}}dt)^2 = \kappa_{xx}dx^2 + \kappa_{yy}dy^2 + \kappa_{zz}dz^2 - \kappa_{tt}dt^2 = ds^2$
 $\kappa_{xx} = \kappa_{yy} = \kappa_{zz} = \kappa_{tt} = 1$ is flat space, Minkowski, as in his Dirac equation(1).

(3) Newpde: $\gamma^\mu \sqrt{\kappa_{\mu\mu}} \partial\psi/\partial x_\mu = (\omega/c)\psi$ for e, ν . So we didn't just drop the $\kappa_{\mu\nu}$ (as is done in ref.1)

(4) Here $\kappa_{oo} = 1 - r_H/r = 1/\kappa_{rr}$, $r_H = (2e^2)(10^{40N})/(mc^2)$. The $N = \dots -1, 0, 1, \dots$ fractal scales (next page)

(5) This Newpde κ_{ij} contains a Mandelbrot set(6) $e^2 10^{40N}$ Nth fractal scale source(fig1) term (from eq.13) that also successfully **unifies theoretical physics**. For example:

For $N = -1$ (i.e., $e^2 X 10^{-40} \equiv Gm_e^2$) κ_{ij} is then by inspection(4) the Schwarzschild metric g_{ij} ; so we just derived General Relativity and the gravity constant G from Quantum Mechanics in one line Wow
For $N = 1$ (so $r < r_c$) Newpde zitterbewegung expansion stage explains the universe expansion (For $r > r_c$ it's not observed, per Schrodinger's 1932 paper.).

For $N = 1$ zitterbewegung harmonic coordinates and Minkowski metric submanifold (after long time expansion) gets the De Sitter ambient metric we observe (D16, 6.2).

For $N = 0$ Newpde $r = r_H$ $2P_{3/2}$ state composite $3e$ is the baryons (QCD not required) and Newpde $r = r_H$ composite e, ν is the 4 Standard electroweak Model Bosons (4 eq.12 rotations \rightarrow appendix A)
for $N = 0$ the higher order Taylor expansion (terms) of $\sqrt{\kappa_{ij}}$ gives the anomalous gyromagnetic ratio and Lamb shift *without* the renormalization and infinities (appendix D3): This is very important
So $\kappa_{\mu\nu}$ provides the general covariance of the Newpde. Eq. 4 even provides us space-time r, t .
So we got all physics here by *mere inspection* of this (curved space) Newpde with no gauges!
We fixed it.

So where does that Newpde come from that fixed it? All mathematicians know that the real numbers (ie .rationals & irrationals) can be constructed from Cauchy completeness i.e. real# sets as rational Cauchy sequence limits. So all we did here is show we postulated real#0 by using it to derive a associated rational Cauchy sequence. We did this because that same postulate (of real#0) math *also* implies fundamental theoretical physics. See below "Results" & "Summary".

Recall $1 \equiv 1+0$ and (the list) $0 \equiv 0X0, 1 \equiv 1X1$ defined as $\mathbf{z} = \mathbf{z}\mathbf{z}$:
the simplest algebraic definition of 0 and 1 So we hypothesize:

Postulate *real #0* (so *real 1*) if $\underline{z}'=0$ (and $\underline{z}'=1$) is substituted (plugged) into $z'=z'z'+C$ **eq1** results in *some* $C=0$ constant (ie $\delta C=0$). Thus

• **Plug** in $\underline{z}=0=z_0=z'$ in **eq1**. To find *all* C substitute z' on left (**eq1**) into right $z'z'$ repeatedly and get iteration $z_{N+1}=z_N z_N - C$. Constraint $\delta C=0$ requires we reject the C s for which $-\delta C = \delta(z_{N+1} - z_N z_N) = \delta(\infty - \infty) \neq 0$. The C s that are left over define the **Mandelbrot set** $C_M = C$ with a subset $C=0$, fractal scales $\delta z' = 10^{40N} \delta z$, $N = \text{integer}$. These fractal scales having their own δz then perturb that $\underline{z}=1$ on its own fractal scale so put $z=1+\delta z$ in **eq.1** to get
$$\delta z + \delta z \delta z = C \quad (3)$$

Define $N \leq 0$ as 'observable' fractal scales. Thus define the 'observer' fractal scales as $N \geq 1$ implying C and δz are big in eq.3 so $|\delta z| \gg 1$. Then solve equation 3 as a quadratic equation so
$$\delta z = (-1 \pm \sqrt{1 + 4C}) / 2 = dr + idt \text{ if } C \leq -1/4 \text{ (complex)} \quad (4)$$

Mandelbrot set iteration (i.e., $z_{N+1} = z_N z_N - C$) for this $\delta C = 0$ *extremum* $C = -1/4$ is a rational number Cauchy sequence $-1/4, -3/16, -55/256, \dots, 0$ thereby proving the hypothesis of our above postulated *real#0* math (and **real 1** since $\text{real#1} = 1 + 0 = 1 \cup \text{real0}$)

• **Plug** in $\underline{z}=1$ in $z'=1+\delta z$ in **eq1**, So $\delta C=0 = (\text{eq1 implies eq3}) = \delta(\delta z + \delta z \delta z) = \delta \delta z (1) + \delta \delta z (\delta z) + (\delta z) \delta \delta z = (\text{observer } |\delta z| \gg 1) \approx \delta(\delta z \delta z) = 0 = (\text{plug in eq.4}) = \delta[(dr + idt)(dr + idt)] =$

$$\delta[(dr^2 - dt^2) + i(dr dt + dt dr)] = 0 \quad (5)$$

$= 2D \delta[(\text{Minkowski metric, } c=1) + i(\text{Clifford algebra} \rightarrow \text{eq.7a})] \quad (\equiv \text{Dirac eq})$
Factor **real** eq.5 $\delta(dr^2 - dt^2) = \delta[(dr + dt)(dr - dt)] = 0 = [[\delta(dr + dt)](dr - dt)] + [(dr + dt)[\delta(dr - dt)]] = 0 \quad (6)$
so $-dr + dt = ds, -dr - dt = ds \equiv ds_1 (\rightarrow \pm e)$ Squaring & eq.5 gives circle in e, v (dr, dt) 2nd, 3rd quadrants (7)
& $dr + dt = ds, dr - dt = ds, dr \pm dt = 0$, light cone $(\rightarrow v, \bar{v})$ in same (dr, dt) plane 1st, 4th quadrants (8)
& $dr + dt = 0, dr - dt = 0$ so $dr = dt = 0$ defines vacuum (while eq.4 derives space-time) (9)

Those quadrants give *positive* scalar $dr dt$ in eq.7 (if *not* vacuum) so imply the eq.5 *non* infinite extremum **imaginary** $\equiv dr dt + dt dr = 0 \equiv \gamma^i dr^i dt + \gamma^j dt^j dr = (\gamma^i \gamma^j + \gamma^j \gamma^i) dr dt$ so $(\gamma^i \gamma^j + \gamma^j \gamma^i) = 0, i \neq j$ (from **real** eq5 $\gamma^i \gamma^i = 1$) (7a) Thus from eqs 5, 7a: $ds^2 = dr^2 - dt^2 = (\gamma^i dr + i \gamma^j dt)^2$ Note how eq5 and C_M just fall (pop) out of eq.1, amazing! (These quadrants in e, v plane are used to *illustrate* the W^+, W^-, Z_0, γ 4 Boson SM 4 rotation extreme math of below perturbed eq.7 which is eq.12)

• **Both** $\underline{z}=0, \underline{z}=1$ together (in **eq1**. Use orthogonality to get (2D+2D curved space)). Thus $(z=1) + (z=0) = (dx_1 + idx_2) + (dx_3 + idx_4) \equiv dr + idt$ given $dr^2 - dt^2 = (\gamma^i dr + i \gamma^j dt)^2$ if $dr^2 \equiv dx^2 + dy^2 + dz^2$ (3D orthogonality) so that $\gamma^i dr \equiv \gamma^x dx + \gamma^y dy + \gamma^z dz, \gamma^i \gamma^i + \gamma^j \gamma^j = 0, i \neq j, (\gamma^i)^2 = 1$, rewritten (κ_{ii} from $N=0$ C_M perturbation of $N=1$, eqs 7, 13-15) as $(\gamma^x \sqrt{\kappa_{xx}} dx + \gamma^y \sqrt{\kappa_{yy}} dy + \gamma^z \sqrt{\kappa_{zz}} dz + \gamma^t \sqrt{\kappa_{tt}} idt)^2 = \kappa_{xx} dx^2 + \kappa_{yy} dy^2 + \kappa_{zz} dz^2 - \kappa_{tt} dt^2 = ds^2$. Multiply both sides by $1/ds^2$ and $\delta z^2 \equiv \psi^2$ use eq.11 circle result $i \partial \delta z / \partial r = (dr/ds) \delta z$ inside brackets () get 4D QM $\gamma^\mu (\sqrt{\kappa_{\mu\mu}}) \partial \psi / \partial x_\mu = (\omega/c) \psi \equiv \text{Newpde}$ for $e, v, \kappa_{00} = 1 - r_H/r = 1/\kappa_{rr}, r_H = e^2 X 10^{40N}/m$ ($N = -1, 0, 1, \dots$). So $\kappa_{\mu\nu}$ carries the general covariance (eq.13-15) and **Postulate 1** $\rightarrow \text{Newpde}$

Results: of (merely plugging $\underline{z}'=0, \underline{z}'=1$ into **eq.1**) **postulate 1:** (1) backups: davidmaker.com **Newpde:** $N=0$, stable $r=r_H$ composite (part II) $3e 2P_{3/2}$ is baryons (QCD not required), SM is the extreme of 4 e, v quadrant rotations. $N=-1$ is GR. Expansion stage of $N=1$ cosmological sfractal cale $\delta z' = \delta z e^{i\omega t}$ Dirac eq zitterbewegung oscillation is the cosmological expansion, for $N=0$ the 3rd order Taylor expansion component (1) of $\sqrt{\kappa_{rr}}$ gets the anomalous gyromagnetic ratio so don't need the renormalization infinities. So we get the physics here.

Math: We use that $1+c \equiv 1 \cup c$ to define above *list-define* (ring-field) algebra and note again that iteration gives a Cauchy sequence limit of real# eigenvalues, so we get the rel# math as well with no new axioms.

Thus (with the math&physics) we understand *everything* (eg GR, cosmology, QM,e,v SM, baryons, rel#).

•So the *simplest idea imaginable 1* implies all *fundamental math-physics*. no more, no less (eg., We simply have 4D and not a myriad of other dimensions)

Conclusion: So by merely (plugging 0,1 into eq.1) **postulating 1**, out pops the universe, BOOM! easily the most important discovery ever made or that will ever be made again. We finally figured it out.

Reminder: *Real#0* math postulates literally nothing(0) (except **real1** since $1=1+0 \equiv 1 \cup \text{real}\#0$.) The algebraic definition of **1** (and 0) is $z=zz$ (note $z=0,1$) if $C=0$ in the below definition:

Summary: This

Theory is **1** The rest is a (rel#1) definition.

Theory	Real# 1 definition
Postulate 1	is defined algebraically if $z=1$ and $z=0$ (<u>plugged</u>) into $z=zz+C$ eq1 gives some $C=0$ constant (ie $\delta C=0$)
	So
	plug ($\delta C=0$ &) $z=0$ into eq1 iteration (to get <i>all</i> C) get 2D (complex) Mandelbrot set $C_M=C$ (fractal scale N) (This iteration also results in a rational Cauchy sequence confirming 0,1 is a real# comes from above 0,1' definition)
	plug ($\delta C=0$ &) $z=1$ into eq1 get 2D Dirac equation ($(N=1) \equiv$ 'observer' perturbing $N=0$ "observables")
	combine both 2D+2D=4D Newpde using $(dx_1+idx_2)_{z=0}+(dx_3+idx_4)_{z=1}=dr+idt$ & dr 3D orthogonalization
	therefore (So we get all of physics and <i>list-define</i> algebra and Real#math(1 such C_M iteration is rational Cauchy)
postulate 1 \rightarrow Newpde	everything that is physical, no more, no less. See backups at davidmaker.com eg., in "introduction" Ultimate Occam's razor postulate 1 so ultimate physics theory, we completely understand the universe.

Backups for (postulate1 \rightarrow Newpde)


I Math Details of postulate1

Recall $1 \equiv 1+0$ and (the *list*) $0 \equiv 0 \times 0, 1 \equiv 1 \times 1$ defined as $z=zz$:

the simplest algebraic definition of 0 and 1. So we hypothesize:

Postulate real #0 and so **re#1** is defined algebraically if $z=1$ and $z=0$ (plugged) into $z=zz+C$ eq1 gives some $C=0$ constant (ie $\delta C=0$). So

•Plug in $z=0=z_0=z'$ in eq1. To find *all* C substitute z' on left (eq1) into right $z'z'$ repeatedly and get iteration $z_{N+1}=z_N z_N - C$. Constraint $\delta C=0$ requires we reject the Cs for which $-\delta C = \delta(z_{N+1} - z_N z_N) = \delta(\infty - \infty) \neq 0$. The Cs that are left over define the **Mandelbrot set** $C_M=C$ with a subset $C=0$, fractal scales $\delta z' = 10^{40N} \delta z$, $N = \text{integer}$.



These fractal scales having their own δz then perturb that $\underline{z=1}$ on its own fractal scale so substitute ansatz $z=1+\delta z$ in eq.1 to get $(1+\delta z)=(1+\delta z)(1+\delta z)+C$ so that $\delta z+\delta z\delta z=C$ (3) Define $N\leq 0$ as 'observable' fractal scales. Thus define the 'observer' fractal scales as $N\geq 1$ implying (from equation 3) that for the 'observer' $|\delta z|\gg 1$. Then solve equation 3 as a quadratic equation so that $\delta z=(-1\pm\sqrt{1+4C})/2=dr+idt$ if $C<-1/4$ (complex) (4) The Mandelbrot set iteration formula (i.e., $z_{N+1}=z_N z_N - C$) for this $\delta C=0$ extremum $C=-1/4$ is a rational# Cauchy seq. $-1/4, -3/16, -55/256, \dots, 0$ confirming the real#0 Cauchy completeness. Thus also $\mathbf{1}$ in above $\mathbf{1}\equiv 1\cup 0$ is a real# verifying **postulate 1**.

• Plug in $\underline{z=1}$ in $z'=1+\delta z$ in eq1, So $\delta C=0$ (eq1 implies eq3) $=\delta(\delta z+\delta z\delta z)=\delta\delta z(1)+\delta\delta z(\delta z)+(\delta z)\delta\delta z$ (observer $|\delta z|\gg 1$) $\approx\delta(\delta z\delta z)=0$ (plug in eq.4) $=\delta[(dr+idt)(dr+idt)]=\delta[(dr^2-dt^2)+i(dr dt+dt dr)]=0$ (5) $=2D \delta[(\text{Minkowski metric, } c=1)+i(\text{Clifford algebra}\rightarrow\text{eq.7a})]$ (\equiv Dirac eq) Factor real eq.5 $\delta(dr^2-dt^2)=\delta[(dr+dt)(dr-dt)]=0=[\delta(dr+dt)](dr-dt)+[(dr+dt)\delta(dr-dt)]=0$ (6) so $-dr+dt=ds, -dr-dt=ds\equiv ds_1(\rightarrow\pm e)$ Squaring&eq.5 gives circle.in e, v (dr,dt) 2nd,3rdquadrants (7) & $dr+dt=ds, dr-dt=ds, dr\pm dt=0$, light cone ($\rightarrow v, \bar{v}$) in same (dr,dt) plane 1st,4thquadrants (8) & $dr+dt=0, dr-dt=0$ so $dr=dt=0$ defines vacuum (while eq.4 derives space-time) (9) Those quadrants give positive scalar $dr dt$ in eq.7 (if not vacuum) so imply the eq.5 non infinite extremum imaginary $\equiv dr dt+dt dr=0\equiv\gamma^i dr^j dt + \gamma^j dt^i dr = (\gamma^i \gamma^j + \gamma^j \gamma^i) dr dt$ so $(\gamma^i \gamma^j + \gamma^j \gamma^i)=0, i\neq j$ (from real eq5 $\gamma^i \gamma^i=1$) (7a)

Thus from eqs5,7a: $ds^2 = dr^2 - dt^2 = (\gamma^r dr + i\gamma^t dt)^2$ Note how eq5 and the Mandelbrot set just fall (pop) out of eq.1, amazing! (These quadrants in that e, v plane are used merely to illustrate the 4 Boson(i.e., W_+, W_-, Z_0, γ) SM 4 rotation extreme of perturbed eq.7 which is eq.12.)

We square eqs.7 or 8 or 9 $ds_1^2 = (dr+dt)(dr+dt) = (-dr-dt)(-dr-dt) = [dr^2+dt^2] + (dr dt+dt dr) \equiv ds^2+ds_3=ds_1^2$. Circle $\equiv \delta z = dse^{i\theta} = dse^{i(\Delta\theta+\theta_0)} = dse^{i((\cos\theta dr + \sin\theta dt)/(ds)+\theta_0)}$, $\theta_0=45^\circ$ (δz in fig.7). We define $k\equiv dr/ds, \omega\equiv dt/ds, \sin\theta\equiv r, \cos\theta\equiv t, dse^{i45^\circ}\equiv ds'$. Take ordinary derivative dr (since flat space)

$$\text{of 'Circle' } \frac{\partial \left(dse^{i\left(\frac{rdr}{ds} + \frac{tdt}{ds}\right)} \right)}{\partial r} = i \frac{dr}{ds} \delta z \text{ so } \frac{\partial (dse^{i(rk+wt)})}{\partial r} = ik\delta z, \quad k\delta z = -i \frac{\partial \delta z}{\partial r} \quad (11).$$

(So given $\delta z\equiv\psi, F\equiv k$ then from eq.11 $\langle F \rangle^* = \int (F\psi)^* \psi d\tau = \int \psi^* F\psi d\tau = \langle F \rangle$. Therefore k is Hermitian). Also from right side real# Cauchy seq. starting at $-1/4$ rational #iteration, is the same as the the Mandelbrot set iteration(7), Ch.2,sect.2, with small C 0 =limit making real eigenvalues (eg.,noise) likely. Thus the Mandelbrot set iteration here did double duty also as proof of the real number eigenvalues in eq.11. The observables $dr\rightarrow k\rightarrow p_r$ condition gotten from eq.11 operator formalism(10) thereby converts eq.7-9 into Dirac eq. pdes (4XCircle extreme in left side fig.1 thereby implies circle observability eq11 which we can then pull out of the zoom.

Note this is then the $N=0$ curved space δz in eq12 allowing us to define $N=0$ as the “observables” fractal scale and $N=1$ as the “observer” scale with its eq5 flat space instead so with no ‘observables’ to observe). Cancel that e^{i45° coefficient ($45^\circ=\pi/4$) then multiply both sides of eq.11 by \hbar and define $\delta z \equiv \psi$, $p \equiv \hbar k$. Eq.11: the familiar ‘observables’ p_r in $p_r \psi = i \hbar \frac{\partial \psi}{\partial r}$ (11) Repeat eq.3 for the τ , μ respective δz lobes in fig.6 so they each have their own neutrino ν : Lepton generations

$\delta C=0$ Extremum on Circle 4X sequence shapes (fig1) In Mandelbrot set pulls it out of zoom clutter because of the above 4X circle observability sequence in fig1

$\delta C=0$ as usual applies to a differential extremum $\delta C = \Sigma(\partial C/\partial x_i) dx_i$ and we must in its final application apply it to $N \leq 0$ observables $C \approx \delta z$ (otherwise why bother?). So $\delta C = \left(\frac{\partial C}{\partial r}\right)_t dr + \left(\frac{\partial C}{\partial t}\right)_r idt = 0$. So for that fig.1 4X sequence of circles $dr dt = darea_M \neq 0$ (so eq.11 observables) the real $\delta C=0$ extremum given the decreasing circle radius sequence $\lim_{m \rightarrow \infty} \frac{\partial C}{\partial area_m} dr_m = KX0 = 0$ (since $dr_\infty \approx 0$) at Feigenbaum point $= f^\alpha = (-1.40115, i0) = C_M \equiv \text{end}$ and is the *ultimate realization of $\delta C=0$* . So random circles in the zoom don’t do $\delta C=0$. Note if a circle (or many circles) is rotated (U), translated (D), shrunk (S) equally in both dimensions (i.e., $(\partial x^j/\partial x^k)^{\beta j} = f^k \equiv \begin{bmatrix} f_{1N} \\ f_{2N} \end{bmatrix} = S_N \begin{bmatrix} U_{11} & U_{12} \\ U_{21} & U_{22} \end{bmatrix} \begin{bmatrix} f \\ 0 \end{bmatrix} + \begin{bmatrix} D_{1N} \\ D_{2N} \end{bmatrix}$) it is still a circle, eq.11 still holds, so *it’s still an observable* as seen in the N fractal scale zoom. Thus you can pick out from that zoom these fig.1 Mandelbrot set extremum 4Xdiameter circles as the only **observables** and $\delta C=0$ extremum geometry in all that clutter. Reset the zoom, restart at such $S_N C_M = 10^{40N} C_M$ in eq.13.

Real eq.5 implies Minkowski metric and so Lorentz transformation boosts γ on scale N to get the small C of postulate 1.

For $N=0$ observable Postulate 1 also implies a small C in eq.1 which implies a eq.5 Lorentz contraction (9) $1/\gamma$ boosted frame of reference (fig.6) in $N=0$ eq.3 **small $C=C_M/\gamma \equiv C_M/\xi_1 = \delta z'$** $z=1+\delta z$ and $\delta C_M = (\delta \xi) \delta z + \xi \delta \delta z = 0$. So must add $N=0$ curved space perturbation $\delta z'$ in eqs.11,12

for $z=1$ δz is small so $\delta \xi$ and ξ can be large (**unstable large mass $\tau + \mu$** , sectD4). (11a)

for $z=0$ $|\delta z|$ is large so $\delta \xi$ and ξ can be small (**stable small mass: electron** ground state δz (11b))

For $N=1$ $\delta z = dr$ gets small relative to 1 at high energy Lorentz boost δz but still keeps **$dr^2 - dt^2 = ds^2$** constant so merely results in slightly modified eq.7: $(dr - \delta z') + (dt + \delta z') \equiv dr' + dt' = ds$ (12)

since ds must remain a constant implying angle perturbation from $\theta_0 = 45^\circ$ on the above **ds Circle**

For $N_{ob}=0$ (observer at $N=1$) and eq. 7 $dr + dt = ds$ the r, t axis’ are the max extremum for ds^2 , and the ds^2 at 45° is the min extremum ds^2 so each $\Delta \theta = \pm 45^\circ$ is pinned to an axis’ so extreme $\Delta \theta \approx \pm 45^\circ = \delta z'$. So in eq.12 the 4 rotations $45^\circ + 45^\circ = 90^\circ$ define 4 Bosons (see **appendix A**). But

for $N=-1$ $45^\circ - 45^\circ = 0^\circ$ $N_{ob} < 0$ then contributes so you also have other (smaller and **infinitesimal $N=-1$**) fractal scale extreme $\delta z'$ (eg., tiny Feigenbaum pts so $N=1$ $dr=r$, for $N_{ob}=-1$) so metric coefficient

$\kappa_{rr} \equiv (dr/dr')^2 = (dr/(dr - (C_M/\xi_1)))^2 = 1/(1 - r_H/r)^2 = A_1/(1 - r_H/r) + A_2/(1 - r_H/r)^2$. The partial fractions A_I can be split off from RN and so

$$\kappa_{rr} \approx 1/[1 - ((C_M/\xi_1)r)] \tag{13}$$

(C_M defined to be e^2 charge, $\gamma \equiv \xi_1$ mass). So: $ds^2 = \kappa_{rr} dr'^2 + \kappa_{oo} dt'^2$ (14)

From eq.7a $dr' dt' = \sqrt{\kappa_{rr}} dr' \sqrt{\kappa_{oo}} dt' = dr dt$ so $\kappa_{rr} = 1/\kappa_{oo}$ (15)

We can then do a rotational dyadic coordinate transformation of $\kappa_{\mu\nu}$ to get the Kerr metric which is all we need for our applications(9). Recall also from eqs5,7a that $dr^2-dt^2=(\gamma^r dr+i\gamma^t dt)^2$

Both $z=0, z=1$ together using orthogonality to get (2D+2Dcurved space). Thus $(z=1)+(z=0)=(dx_1+idx_2)+(dx_3+idx_4)\equiv dr+idt$ given $dr^2-dt^2=(\gamma^r dr+i\gamma^t dt)^2$ if $dr^2\equiv dx^2+dy^2+dz^2$ (3D orthogonality) so that $\gamma^r dr\equiv \gamma^x dx+\gamma^y dy+\gamma^z dz$, $\gamma^i \gamma^j+\gamma^j \gamma^i=0$, $i\neq j, (\gamma^i)^2=1$ (B2), rewritten (with eq14)

$(\gamma^x \sqrt{\kappa_{xx}} dx+\gamma^y \sqrt{\kappa_{yy}} dy+\gamma^z \sqrt{\kappa_{zz}} dz+\gamma^t \sqrt{\kappa_{tt}} idt)^2=\kappa_{xx} dx^2+\kappa_{yy} dy^2+\kappa_{zz} dz^2-\kappa_{tt} dt^2=ds^2$. Multiply both sides by $1/ds^2$ & $(\delta z/\sqrt{dV})^2\equiv \psi^2$ and using operator eq 11 inside the brackets () get **Newpde**

$\gamma^\mu (\sqrt{\kappa_{\mu\mu}}) \partial\psi/\partial x_\mu=(\omega/c)\psi$ for e, ν , $\kappa_{00}=1-r_H/r=1/\kappa_{rr}$ $r_H=e^2 X 10^{40N}/m$ ($N=. -1, 0, 1, .$) (16)
 $=C_M/\xi_1$ (from* eq.13) $C_M=Fiegenbaum$ point. Also $C_M/\xi=r_H=$

*small C so big $\xi=\gamma$ boost so $z=zz$ so **postulate 1**. So we really did just postulate 1. So

Postulate 1 \rightarrow **Newpde**

* C_M/ξ_1 is ξ small C boost for $z=zz$ so postulate1 from Newpde $r=r_H$ $2P_{3/2}$ stable state. See fig6. The 4 eq.12 Newpde e, ν rotations at $r=r_H$ are the 4 W^+, γ, W^-, Z_0 SM Bosons (appendixA).

So Penrose's intuition(6) was right on! There *is* physics in the Mandelbrot set, all of it.

2.1 Newpde Oscillation of $\delta z(\equiv\psi)$ on $N\geq 1$ fractal scale is Cosmology

From Newpde eq16 (eg., eq.1.13 Bjorken and Drell) $i\hbar \frac{\partial\psi}{\partial t} = \frac{\hbar c}{i} \left(\alpha_1 \frac{\partial\psi}{\partial x^1} + \alpha_2 \frac{\partial\psi}{\partial x^2} + \alpha_3 \frac{\partial\psi}{\partial x^3} \right) +$

$\beta mc^2 \psi = H\psi$. For electron at rest: $i\hbar \frac{\partial\psi}{\partial t} = \beta mc^2 \psi$ so: $\delta z = \psi_r = w^r(0) e^{-i\varepsilon_r \frac{mc^2}{\hbar} t}$ $\varepsilon_r=+1,$

$r=1, 2; \varepsilon_r=-1, r=3, 4$): This implies an oscillation frequency of $\omega=mc^2/\hbar$. which is fractal here. ($\omega=\omega_0 10^{-40N}$). So the eq.12 the 45° line has this ω oscillation as a (that eq.7-9 δz variation) rotation. On our own fractal cosmological scale we are in the expansion stage of one such oscillation. Thus the fractalness of the Newpde explains cosmology. The next higher cosmological scale is independent (but still connected by superposition of speeds implying a separation of variables result: $i\hbar \frac{\partial\psi}{\partial t} = \beta \sum_N (10^{-40N} (\omega t)_{\varepsilon+\Delta\varepsilon}) \psi = \beta \sum_N (10^{-40N} m_{\varepsilon+\Delta\varepsilon} c^2 / \hbar) \psi$). By the way fractal scale $N=1$ the 45° small Mandelbulb chord ε (Fig6) is now, given this ω , getting larger with time so $1-t \propto \varepsilon$. But the tauon 68.74° is stationary so its mass can be set to 1. So at this time (relative to the tauon) the muon $=\varepsilon=.06$, electron $\Delta\varepsilon=.0005899$. So

cosmologically for stationary $N=2$ $\delta z=\sqrt{\kappa_{00}} dt=e^{-i\varepsilon_r \frac{mc^2}{\hbar} t} \rightarrow e^{i(\varepsilon+\Delta\varepsilon)}$ (17)

But seen from inside at $N=1$ (D18) $E=1/\sqrt{\kappa_{00}}=1/\sqrt{(1-r_H/r)}$ then $r<r_H$ & E becomes imaginary in $e^{iEt/\hbar}=\delta z=\sqrt{\kappa_{00}} dt= e^{-i\varepsilon_r \frac{mc^2}{\hbar} t} \rightarrow e^{(\varepsilon+\Delta\varepsilon)}$ (17a)

This $N=0$ and $N=-1$ δz is the source of the small rotation in eq.12. Later we see that $N=0$ high energy scattering drives the $\delta\delta z$ term (/ds) to the big $\Delta 45^\circ$ extreme (so preferred) jumps (appendixA).

2.2 ambient metric ε (inertial frame dragging reduction) inputs. Eq.D9 is ambient metric which means $N=1$ observer for these ε masses

Postulate 1 (observable) requires that $C\approx 0$ in equation 1. Note also that the **real** component of eq.5 is the Minkowski metric implying these γ boosts. Recall eq.3 $\delta z+\delta z\delta z=C$. So for $N=1$ observer $|\delta z|\gg 1$ so $\delta z\delta z=C$. Given eq.3 for $N=0$ $|\delta z|\gg |\delta z\delta z|$, $C\approx\delta z$ sect.1 for $N=0$. Note also our above circle e electron -dr $\Delta\varepsilon$ intersection ground state -dr is at 45° (2nd & 3rd quadrants) for minimum ds^2 . So following the energy increase for Newpde states μ then is not a constant in time because of $N=1$ eq.12 angle Newpde zitterbewegung variable time contribution (eq.17) to

the δz chord perturbation of the 45° (fig6 below). For next higher energy the $68.7^\circ = \text{Arctan}(\delta z/C_M)$ is from eq.4 quadratic equation solution at the Feigenbaum point.(so it gives our *2fundamental* excited state Mandelbulb) mass τ that does not change over cosmological time in $N=1$ allowing us to normalize it to 1). Note these are identical to eq.7-9 of the section 1 eq.3 application for the τ , μ respective δz lobes in fig.6 so they each have their own neutrino v.eq.7,8,9 with its electron' and neutrino still the core equations even for the muon and tauon thereby deriving the 3 generations of leptons.

Stability of composite 3e: (Newpse stable $2P_{3/2}$ at $r=r_H$ state)

We can actually calculate m_p from the quantization of the magnetic flux $h/2e=\Phi_0=BA$ (partII) using the Newpde ground state $z=0$ three electron (S_1, S_2, S_3), $e=e+e-e$ states of the Newpde with LS coupling minimal energy ($\mathbf{J}=\mathbf{L}+\mathbf{S}=1-\frac{1}{2}-\frac{1}{2}+\frac{1}{2}=\frac{1}{2}$) with two orbiting relativistic positrons γm_e for $2P_{3/2}$ at $r=r_H$, so $3e=(\gamma m_e+\gamma m_e)=m_p$ Stability is implied by $(dt')^2=(1-r_H/r)dt^2$ since clocks stop ($dt'=0$) at $r=r_H$. That 3rd mass also reverses the pair annihilation with virtual pair creation inside the r_H 2D area given $\sigma=\pi r_H^2 \approx (1/20)$ barns which is the reason why only composite 3e or its multiples gives stability.

Note these 2D τ, μ Mandelbulbs can be on a flat 2D ($z=1$) or this spherical 2D shell ($z=0$) That makes this spherical shell at $r=r_H$ the only other stable 2D space (in addition to these $z=1$ flat 2D) Newpde ground state to define these Mandelbulbs on. Thus high energy 2D $\tau+\mu$ Mandelbulbs provide 3e stability in μ and 3e in τ so $\mu+\tau=3e+3e=(\gamma m_e+\gamma m_e)_\tau+(\gamma m_e+\gamma m_e)_\mu$ as 2 $2P_{3/2}$ orbitals with S and L inside the horizon r_H so unobserved so all that is seen from the outside is (no longer the inside 2P) net $\mathbf{J}=\mathbf{S}'=\frac{1}{2}$.

For $N=0$ observable

$z=0$, $r=r_H$ 11b, the high energy $r=r_H$ 2D spherical shell then is a domain of these same 2D Mandelbulbs μ , τ giving on the 2D shell: $\mu+\tau=3e+3e=(\gamma m_e+\gamma m_e)_\tau+(\gamma m_e+\gamma m_e)_\mu=3e+3e=m_p+m_p$. two body motion equipartition of energy of the interacting positrons in each of two **baryons** each with $\mathbf{J}=\mathbf{S}'=\frac{1}{2}$. Eq 11b so for each positron $\delta z'=r_H=C_M/\xi_o=C_M/m_e$ in eq.12.

$z=1$, 11a, $r'_H \ll r_H$ (so not that shell) because for $z=1$ $\xi_1 \gg \xi_o$ $\lambda=h/mc$ =Compton wavelength, $2\pi r'_H=\lambda$, $m=\xi_1$. Again 3e for each of 2D free space domain high energy quasi stable μ, τ ,: $\tau+\mu=3e+3e=2$ free space **leptons** each with $\mathbf{J}=\mathbf{S}'=\frac{1}{2}$. **11a** so $\delta z=r'_H=C_M/\xi_1=C_M/(\tau+\mu)$ (18) in eq12

For $N=1$ observer eq.3 implies $C=\delta z \delta z/\xi$ so that $\xi=C/\delta z \delta z=C/(\text{Mandelbulb radius})^2=\text{mass}$ (from fig.6). or as a fraction of τ , with $2m_p=\tau+\mu+e=\xi_1$ electron $\Delta \epsilon=.00058$ (19)

Postulate 1 implied finally

But γ (observer) $=\gamma$ (observable) so for the $N=0$ observable we got the γ from the $N=1$ observer case in $r_H=C_M/\gamma=C_M/\xi=C$ for small C and so postulate 1. Thus we really did just **postulate 1**.




Fig.6 Conclusion. So the small C at the end was required. So we really did just postulate 1

So we just do *what is simplest* (let Occam be your guide), just **postulate 1**: the physics (Newpde) will then follow, top down:

*** Ultimate Occam's Razor (observable)**

It means here *ultimate* simplicity, the *simplest* idea imaginable. So for example $z=zz$ is *simpler* than $z=zzzz$. Therefore **1** in this context (uniquely algebraically defined by $z=zz$) is this ultimate Occam's razor **postulate** since 0 (also from $z=zz$) postulates literally *nothing*.

But postulate real 0 is what initially comes out that postulate math so that $1 \equiv 1 \cup \text{real } 0$ implies **postulate 1**.

2.3 Left end small drdt (eq.6) extremum Fiegenbaum point Fractalness

The Fiegenbaum point (11a) is the only part of the Mandelbrot set we use. At the Fiegenbaum point (imaginary) time $X10^{-40} = \Delta$ and real -1.40115 . Since $|C_M| \gg 0$ in eq.2 postulated eq.1 $z=zz$ implies a boosted SR Lorentz transformation universal reference frame to random (since this transformation cancels noise C in eq.2, fig6), small C_M subset $C \approx \delta z'$ (from eq.3) = real distance = $\text{real} \delta z / \gamma = 1.4011 / \gamma = C_M / \gamma \equiv C_M / \xi_1$ using large ξ_1 . Note at the Fiegenbaum point distance $1.4011 / \gamma$ shrinks a lot but time $X10^{-40} \gamma$ doesn't get much bigger since it was so small to begin with at the Fiegenbaum point. Eq.1 then means we have Occam's razor optimized **postulated 1**. Given the New pde r_H we only see the $r_H = e^{2 \cdot 10^{40N}} / m$ sources from our N=0 observer baseline. We never see the $r < r_H$ <http://www.youtube.com/watch?v=0jGaio87u3A> which explores the Mandelbrot set interior near the Fiegenbaum point. Reset the zoom start at such extremum $S_N C_M = 10^{40N} C_M$ in eq.13. The splits are in 3 directions from the orbs. There appear to be about 2.5 splits going by each second (given my PC baud rate) and the next Mandelbrot set comes up in about 62 seconds. So $3^{2.7 \times 62} = 10^N$ so $172 \log 3 = N = 82$. So there are 10^{82} splits. So there are about 10^{82} splits per initial split. But each of these Mandelbrot set Fiegenbaum points is a $C_M / \xi \equiv r_H$ in electron (eq.13 above). So for each larger electron there are **10^{82} constituent electrons**. Also the scale difference between Mandelbrot sets as seen in the zoom is about 10^{40} , **the scale change** between the classical electron radius and 10^{11} ly with the C noising giving us our fractal universe.

Recall again we got from eq.3 $\delta z + \delta z \delta z = C$ with quadratic equation result:


$$\delta z = \frac{-1 \pm \sqrt{1-4C}}{2}$$

is real for noise $C < 1/4$ creating our noise on the N=0 th fractal scale. So $1/4 = (3/2)kT / (m_p c^2)$. So T is 20MK. So here we have *derived the average temperature of the universe* (stellar average). That $z' = 1 + \delta z$ substitution also introduces Lorentz transformation

rotational and translation noise that does not effect the number of splits, analogous to how a homeomorphism does not change the number of holes (which is a Topological invariant). So the excess C noise (due to that small C' boost) causes the Feigenbaum point neighborhood internal structure to become randomized (as our present universe is) but the number of electrons (10^{82}) remains invariant. See appendix D mixed state case2 for further organizational effects. $N=r^D$. So the **fractal dimension**= $D=\log N/\log r=\log(\text{splits})/\log(\#r_H \text{ in scale jump})$
 $=\log 10^{80}/\log 10^{40}=\log(10^{40})^2/\log(10^{40})=2$. (See appendix E for Hausdorff dimension & measure) which is the same as the 2D of eq.4 and the Mandelbrot set. The next smaller (subatomic) fractal scale $r_1=r_H=2e^2/m_e c^2$, $N=0$ th, $r_2=r_H=2GM/c^2$ is defined as the $N=1$ th where $M=10^{82}m_e$ with $r_2=10^{40}r_1$ So the Feigenbaum pt. gave us a lot of physics:
 eg. **#of electrons in the universe, the universe size, temp.**

Iteration Math

Mandelbrot set iteration sequence z_n $C_M=-1/4$, $z_0=0$ same as Cauchy seq. since it begins with rational number $-1/4$, allowing the (C' uncertainty) dr neighborhood of 0 to have a nonzero probability of being a real number and thereby giving real eigenvalues to the equation 11 operator formalism after the small C' boost to get observability around $dr=0$. $dr=0$.
 So $\delta z \approx \text{zero}$ ($N=0$ fractal scale) is a real number which makes the $z=1$ in $z=1+\delta z \approx 1+0$ a real number thereby **confirming our original postulate real #1**. The postulate 1 also gives the *list-define* math (B2) *list* cases $1 \cup 1 = 1+1 = 2$, *define* $a=b+c$ (So no other math axioms but 1.)
 That means the **mathematics and the physics** come from (**postulate 1** \rightarrow **Newpde**): *everything*. Recall from eq.7 that $dr+dt=ds$. So combining in quadrature eqs 7&11 $SNR\delta z=(dr/ds+dt/ds)\delta z=((dr+dt)/ds)\delta z=(1)\delta z$ (11c,append) and so having come *full circle* back to sect.1 postulate 1 as a real eigenvalue (1 \equiv **Newpde** electron). So, having come *full circle* then: (**postulate 1** \Leftrightarrow **Newpde**), back to our section 1. So we rewrite our title:
 “The Ultimate Occam’s razor theory (ie 1) is *the same as* the ultimate math-physics theory (ie **Newpde**)”.



2.4 Results: Recall from ultimate Occam’s razor **Postulate 1** we got the Newpde. We note in reference 5 on the first page that we also get the *actual* physics with the Newpde. Thus the usual postulating of hundreds of Lagrange densities(fig.11), free parameters, dimensions, etc., is senseless.

For example (appendixC) **Newpde composite 3e** $2P_{3/2}$ at $r=r_H$ is the proton: That B flux quantization(C3) implies a big proton mass implying 2 high speed $\gamma=917$ positrons and so the Fitzgerald **contracted E field lines are the strong force**: we finally understand the strong force! (bye,bye QCD). So these *two* positrons then have big mass *two body* motion(partII) so also **ortho(s,c,b) and para(t) Paschen Back excited (hadron multiplet) states** understood (partII) $N=0$ extreme perturbation rotations of $N=1$ eq.12 implies **Composite e, ν** at $r=r_H$ giving **the**

electroweak SM (appendixA) **Special relativity** is that eq.5 Minkowski result. **With the Eqs.16 Newpde ψ** (appendix C) **we finally understand Quantum Mechanics** for the first time and eq.4 gave us a **first principles derivation of r,t space-time** for the first time. That Newpde $\kappa_{\mu\nu}$ metric (In eq.14), on the $N=-1$ next smaller fractal scale(1) so $r_H=10^{-40}2e^2/m_e c^2 \equiv 2Gm_e/c^2$, is the Schwarzschild metric since $\kappa_{00}=1-r_H/r=1/\kappa_{rr}$ (15): we **just derived General Relativity(gravity) from quantum mechanics in one line**. The Newpde zitterbewegung expansion component ($r < r_C$) on the next larger fractal scale ($N=1$) is the universe expansion sect.2.1: **we just derived the expansion of the universe in one line**. The third order terms in the Taylor expansion of the Newpde $\sqrt{\kappa_{\mu\nu}}$ give those precision QED values (eg.,Lamb shift sect.D) allowing us to **abolish the renormalization and infinities**.

So there is no need for those many SM Lagrangian density postulates (fig11) anymore, just **postulate 1** instead.

Intuitive Notion (of postulate 1 \leftrightarrow Newpde)

The Mandelbrot set introduces that $r_H = C_M/\xi_1$ horizon in $\kappa_{00}=1-r_H/r$ in the Newpde, where C_M is fractal by $10^{40}X$ scale change(fig.2) So we have found (davidmaker.com) that: Given that fractal selfsimilarity astronomers are observing from the inside of what particle physicists are studying from the outside, that **ONE New pde e** electron r_H , **one** thing (fig.1). *Everything* we observe big (cosmological) and small (subatomic) is then that (*New pde*) r_H , even baryons are composite **3e**. So we understand, *everything*. This is the only Occam's razor first principles theory

Summary: So instead of doing the usual powers of 10 simulation we do a single power of 10^{40} simulation and we are immediately back to where we started!




fig2

(↑lowest left corner) Object B caused perturbation structure jumps: void→galaxy→globular,,etc.

References

- (6) Penrose in a utube video implied that the Mandelbrot set might contain physics. Here we merely showed how to find it. For example the (fractal Mandelbulb neighborhood area $|\text{drdt}| > 0$ of the) Feigenbaum point is a subset (containing that 10^{40} X selfsimilar scale jump: Fig1)
- (7) Cantor: Ueber die Ausdehnung eines Satzes aus der Theorie der trigonometrischen Reihen, "Ueber eine elementare Frage der Mannigfaltigkeitslehre" Jahresbericht der Deutschen Mathematiker-Vereinigung." Mandelbrot set iteration sequence z_n $C_M = -1/4$, $z_0 = 0$ same as Cauchy seq. since it begins with rational number $-1/4$, allowing the (C' uncertainty) dr neighborhood of 0 to have a nonzero probability of being a real number and thereby giving real eigenvalues to the equation 11 operator formalism after the small C' boost to get observability around $dr = 0$. $dr = 0$. So $\delta z \approx \text{zero}$ ($N=0$ fractal scale) is a real number which makes the $z=1$ in $z=1+\delta z \approx 1+0$ a real number thereby confirming our original postulate real #1
- (8) Tensor Analysis, Sokolnikoff, John Wiley
- (9) The Principle of Relativity, A Einstein, Dover
- (10) Quantum Mechanics, Merzbacher, John Wiley
- (11) lemniscate circle sequence (Wolfram, Weisstein, Eric)

(12) **appendix A for finite larger $N_{ob} = 0$ required extremum to extremum rotations (jumps) at high interaction COM energies (analogous to a hydrogen atom principle quantum number $N=1$ to $N=2$ jump)**

Recall from sect.1 eq.3 that $\delta C = \delta(\delta z + \delta z \delta z) = \delta \delta \delta z (1) + \delta \delta z (\delta z) + (\delta z) \delta \delta z = \delta C = 0$ so C is split between $\delta \delta z$ noise and $\delta z \delta z$ classical invariance ds^2 proper time.

Recall at $N=0$ the $N=1$ $|\delta z| \gg 1$ & $C_M \gg 1$. So $\delta z \delta z \approx C_M$ there. So equation 5 holds then. But $\frac{\delta z'}{ds} = \pm 45^\circ$ ($\pi/4$) extremum to extremum observable $N=0$ (SM) is also a solution for observer $N=1$ at high interaction COM energies. $N=-1$ is part of the *more general* $N_{ob} < 0$ eq.13-15 case of sect.1 that also allows infinitesimal perturbations.

So for high interaction energies as the γ boosted observer $\delta z/\gamma$, C/γ , gets smaller than the huge $N=1$ scale (so higher energy, smaller wavelength, beam probes) $\delta \delta z(1)/ds$ noise angle gets relatively larger (relative to $\delta(\delta z \delta z)/ds$, sect.1) until finally the next smaller $N=0$ (and next smaller one after that, $N=-1$) is $N=0$ fractal scale in that sect.1 big angle $\pm 45^\circ$ required extremum solution (Recall 'extremum's are our solutions.) $45^\circ = \pi/4 \approx 1 \approx \delta z'/ds$ (observable) = $C_M \text{end}/ds \equiv \theta$ (in equation 12). So here all four $\theta \pm 45^\circ \times 2$ rotations of **Composite e,v** implied by eq.12. So we have the $N=0$ solutions for $\delta z'$ angle perturbation of $N=1$ for big scattering energies. So observer $\gamma = \text{observed } \gamma$

I \rightarrow II, II \rightarrow III, III \rightarrow IV, IV \rightarrow I required extremum to extremum rotations in eq.7-9 plane Give SM Bosons at high interaction COM energies. $N_{ob} = 0$

For $z=0$ $\delta z'$ is big in $z'=1+\delta z$ and so we have again $\pm 45^\circ$ min ds and so two possible 45° rotations so through a total of two quadrants for $\pm \delta z'$ in eq.12. one such rotation around an around a axis (SM) and the other around a diagonal (SC). Note in fig.3 dr, dt is also a rotation. and so has an eq.11 rotation operator observable θ . Thus from equation 11 for (θ) angle rotations $\theta \delta z \equiv (dr/ds) \delta z = -i \partial(\delta z)/\partial r$ for the first 45° rotation. So we got through one Newpde derivative for each 45° rotation. For the next 45° rotation in fig.4 it is then a second derivative $\theta \theta \delta z' = e^{i\theta} e^{i\theta} \delta z = e^{i(2\theta)} \delta z = (dr/ds)((dr/ds) dr') = -i \partial(-i \partial(dr'))/\partial r = -\partial^2(dr')/\partial r^2$ large angle rotation in figure 3. In contrast for $z=1$, $\delta z'$ small so $45^\circ - 45^\circ$ small angle rotation in figure 3 (so then $N=-1$). Do the same with the time t and get for $z=0$ rotation of $45^\circ + 45^\circ$ (fig.4) then $\theta \theta \delta z' = (d^2/dr^2) z' + (d^2/dt^2) \delta z'$ (A1)




fig.3. for $45^\circ-45^\circ$ so two body (e,v) singlet $\Delta S = 1/2 - 1/2 = 0$ component so pairing interaction (sect.4.5). Also ortho $\Delta S = 1/2 + 1/2 = 1$ making 2 body (at $r=r_H$) $S=1$ Bosons and so a field theory. Note we also get these Laplacians characteristic of the Boson field equations by those $45^\circ+45^\circ$ rotations so eq.16 implies Bosons accompany our leptons (given the $\delta z'$), **so these leptons exhibit "force"**.

Newpde $r=r_H, z=0, 45^\circ+45^\circ$ rotation of composites e,v implied by Equation 12

So $z=0$ allows a large C z rotation application from the 4 different axis' max extremum (of eq.16) branch cuts gives the 4 results: Z,+W, photon bosons of the Standard Model. So we have derived the Standard Model of particle physics in this very elegant way (from the four axis'). You are physically at $r=r_H$ if you rotate through the electron quadrants (I, IV).of eq.7-9. So we have large C_M dichotomic 90° rotation to the next Reimann surface of eq.12, eq.A1 $(dr^2+dt^2)z''$ from some initial extremum angle(s) θ . Eq.12 solutions imply complex 2D plane Stern Gerlach dichotomic rotations using eq.A1 thereby using Pauli matrices σ_i algebra, which maps one-to-one to the quaternion algebra. Using eq.12 we start at some initial angle θ and rotate by 90° the noise rotations are: $C=\delta z'' = [e_L, v_L]^T \equiv \delta z'(\uparrow) + \delta z'(\downarrow) \equiv \psi(\uparrow) + \psi(\downarrow)$ has a eq.12 infinitesimal unitary generator $\delta z'' \equiv U = 1 - (i/2)\epsilon n \cdot \sigma$, $n = \theta/\epsilon$ in $ds^2 = U^\dagger U$. But in the limit $n \rightarrow \infty$ we find, using elementary calculus, the result $\exp(-(i/2)\theta \cdot \sigma) = \delta z''$. We can use any axis as a branch cut since all 4 are eq.16 large extremum so for the 2nd rotation we move the branch cut 90° and measure the angle off the next diagonal since Pauli matrix dichotomic rotations are actually axis rotations, leaving our e and v directions the same. In any case $(dr+dt)z''$ in eq.16 can then be replaced by eq.A1 $(dr^2+dt^2+..) \delta z'' = (dr^2+dt^2+..) e^{\text{quaternion A}}$ Bosons because of eq.A1.

A2 Then use eq. 12 and quaternions to rotate $\delta z''$ since the quaternion formulation is isomorphic to the Pauli matrices. $dr' = \delta z_r = \kappa_{rr} dr$ for **Quaternion A** $\kappa_{ii} = e^{iA_i}$.

Appendix A Quaternion ansatz $\kappa_{rr} = e^{iA_r}$ instead of $\kappa_{rr} = (dr/dr')^2$ in eq.14. $N=0$.

A1 for the eq.12: large $\theta = 45^\circ+45^\circ$ rotation (for $N=0$ so large $\delta z' = \theta r_H$). Instead of the equation 13,15 formulation of κ_{ij} for small $\delta z'$ ($z=1$) and large $\theta = 45^\circ+45^\circ$ we use A_r in dr direction with $dr^2 = x^2 + dy^2 + dz^2$. So we can again use 2D (dr,dt) $E = 1/\sqrt{\kappa_{00}} = 1/\sqrt{e^{iA_i}} = e^{-iA/2}$. The 1 is mass energy and the first real component after that in the Taylor expansion is field energy A^2 . For 2 particles together the other particle ϵ negative means r_H is also negative. Since it is $e_1 * e_2 = r_H$. So $1/\kappa_{rr} = 1 + (-\epsilon + r_H/r)$ is \pm and $1 - (-\epsilon + r_H/r)$ 0 charge. (A0)

For baryons with a 3 particle r_H/r may change sign without third particle ϵ changing sign so that at $r=r_H$. Can normalize out the background ϵ in the denominator of $E = (\tau + \epsilon) / \sqrt{(1 + \epsilon + \Delta\epsilon - r_H/r)}$ for

Can normalize out the background ε in the denominator of $E=(\tau+\varepsilon)/\sqrt{(1+\varepsilon+\Delta\varepsilon-r_H/r)}$ for small conserved (constant) energies $1/\sqrt{(1+\varepsilon)}$ and (so $E=(1/\sqrt{(1+x)})=1-x/2+$) large r (so large λ so not on r_H) implies the normalization is:

$E=(\varepsilon+\tau)/\sqrt{((1-\varepsilon/2-\varepsilon/2)/(1\pm\varepsilon/2))}$, $J=0$ para e, ν eq.9.23 π^\pm, π^0 . For large $1/\sqrt{\Delta\varepsilon}$ energies given small $r=r_H$, Here $1+\varepsilon$ is locally constant so can be normalized out as in

$$E=(\varepsilon+\tau)/\sqrt{(1-(\Delta\varepsilon/(1\pm\varepsilon))-r_H/r)}, \text{ for charged if } -, \text{ ortho } e, \nu J=1, W^\pm, Z_0 \quad (11d)$$




fig4

Fig.4 applies to eq.9 $45^\circ+45^\circ=90^\circ$ case: **Bosons**.

A2 These quadrants were defined in eq.7-9 and used in eq.12. The Appendix A4 derivation applies to the far right side figure. Recall from eq.12 $z=0$ result $C_M=45^\circ+45^\circ=90^\circ$, gets Bosons. $45^\circ-45^\circ=$ leptons. The ν in quadrants II(eq.5) and III (eq.9). e in quadrants I (eq.7) and IV (eq.7). Locally normalize out $1+\varepsilon$ (appendix D). For the **composite** e, ν on those required large $z=0$ eq.9 rotations for $C \rightarrow 0$, and for stability $r=r_H$ (eg., for $2P_{1/2}$, $I \rightarrow II$, $III \rightarrow IV$, $IV \rightarrow I$) unless $r_H=0$ ($II \rightarrow III$) Example:

A4 Quadrants IV \rightarrow I rotation eq.A2 $(dr^2+dt^2+..)e^{\text{quaternion } A}$ =rotated through C_M in eq.16.

example C_M in eq.A1 is a 90° CCW rotation from 45° through ν and anti ν

A is the 4 potential. From eq.9b we find after taking logs of both sides that $A_0=1/A_r$ (A2)

Pretending we have a only two i, j quaternions but still use the quaternion rules we first do the r

derivative: From eq. A1 $dr^2\delta z = (\partial^2/\partial r^2)(\exp(iA_r+jA_0)) = (\partial/\partial r)[(i\partial A_r/\partial r + \partial A_0/\partial r)(\exp(iA_r+jA_0))] = \partial/\partial r[(\partial/\partial r)iA_r + (\partial/\partial r)jA_0](\exp(iA_r+jA_0)) + [i\partial A_r/\partial r + j\partial A_0/\partial r]\partial/\partial r(\exp(iA_r+jA_0)) + (i\partial^2 A_r/\partial r^2 + j\partial^2 A_0/\partial r^2)(\exp(iA_r+jA_0)) + [i\partial A_r/\partial r + j\partial A_0/\partial r][i\partial A_r/\partial r + j\partial/\partial r(A_0)] \exp(iA_r+jA_0)$ (A3)

Then do the time derivative second derivative $\partial^2/\partial t^2(\exp(iA_r+jA_0)) = (\partial/\partial t)[(i\partial A_r/\partial t + \partial A_0/\partial t)$

$(\exp(iA_r+jA_0))] = \partial/\partial t[(\partial/\partial t)iA_r + (\partial/\partial t)jA_0](\exp(iA_r+jA_0)) + [i\partial A_r/\partial r + j\partial A_0/\partial t]\partial/\partial r(\exp(iA_r+jA_0)) + (i\partial^2 A_r/\partial t^2 + j\partial^2 A_0/\partial t^2)(\exp(iA_r+jA_0)) + [i\partial A_r/\partial t + j\partial A_0/\partial t][i\partial A_r/\partial t + j\partial/\partial t(A_0)]\exp(iA_r+jA_0)$ (A4)

Adding eq. A2 to eq. A4 to obtain the total D'Alambertian $A_3+A_4=$

$[i\partial^2 A_r/\partial r^2 + i\partial^2 A_r/\partial t^2] + [j\partial^2 A_0/\partial r^2 + j\partial^2 A_0/\partial t^2] + ii(\partial A_r/\partial r)^2 + ij(\partial A_r/\partial r)(\partial A_0/\partial r) + ji(\partial A_0/\partial r)(\partial A_r/\partial r) + jj(\partial A_0/\partial r)^2 + ii(\partial A_r/\partial t)^2 + ij(\partial A_r/\partial t)(\partial A_0/\partial t) + ji(\partial A_0/\partial t)(\partial A_r/\partial t) + jj(\partial A_0/\partial t)^2$.

Since $ii=-1, jj=-1, ij=-ji$ the middle terms cancel leaving $[i\partial^2 A_r/\partial r^2 + i\partial^2 A_r/\partial t^2] +$

$[j\partial^2 A_0/\partial r^2 + j\partial^2 A_0/\partial t^2] + ii(\partial A_r/\partial r)^2 + jj(\partial A_0/\partial r)^2 + ii(\partial A_r/\partial t)^2 + jj(\partial A_0/\partial t)^2$

Plugging in A2 and A4 gives us cross terms $jj(\partial A_0/\partial r)^2 + ii(\partial A_r/\partial t)^2 = jj(\partial(-A_r)/\partial r)^2 + ii(\partial A_r/\partial t)^2$

$=0$. So $jj(\partial A_r/\partial r)^2 = -jj(\partial A_0/\partial t)^2$ or taking the square root: $\partial A_r/\partial r + \partial A_0/\partial t = 0$ (A5)

$i[\partial^2 A_r/\partial r^2 + i\partial^2 A_r/\partial t^2] = 0, j[\partial^2 A_0/\partial r^2 + i\partial^2 A_0/\partial t^2] = 0$ or $\partial^2 A_\mu/\partial r^2 + \partial^2 A_\mu/\partial t^2 + .. = 1$ (A6)

A4 and A5 are **Maxwell's equations** (Lorentz gauge formulation) in free space, if $\mu=1,2,3,4$.

$$\square^2 A_\mu = 1, \quad \square \bullet A_\mu = 0 \quad (A7)$$

The Lorentz gauge is the only gauge hence it is no gauge at all and we have avoided the Maxwell overdeterminism problem (8eq, 6 unknowns E_i, B_i). Must use Newpde 4D orthogonalization here Amplitudes of physical processes in QED in the noncovariant Coulomb gauge coincide with those in the covariant Lorenz gauge. The Aharonov–Bohm effect depends on a line integral of \mathbf{A} around a closed loop, and this integral is not changed by $\mathbf{A} \rightarrow \mathbf{A} + \nabla\psi$ which doesn't change $\mathbf{B} = \nabla \times \mathbf{A}$ either. So formulation in the Lorentz gauge mathematics works so it is no longer a gauge, we are gaugeless.

A5 Other 45°+45° Rotations (Besides above quadrants IV→I)

For the **composite e,ν** on those required large $z=0$ eq.12 rotations for $C \approx 0$, and for stability $r=r_H$ for $2P_{1/2}$ (I→II, III→IV, II→III) unless $r_H=0$ (IV→I) are:

Ist→IInd quadrant rotation is the W^+ at $r=r_H$. Do similar math to A2-A7 math and get instead a Proca equation The limit $\varepsilon \rightarrow 1 = \tau$ (D13) in ξ_1 at $r=r_H$. since Hund's rule implies $\mu = \varepsilon = 1S_{1/2} \leq 2S_{1/2} = \tau = 1$. So the ε is negative in $\Delta\varepsilon/(1-\varepsilon)$ as in case 1 charged as in appendix C1 case 2.

$E = 1/\sqrt{\kappa_{00}} - 1 = [1/\sqrt{(1-\Delta\varepsilon/(1-\varepsilon)-r_H/r)}] - 1 = [1/\sqrt{(\Delta\varepsilon/(1-\varepsilon))}] - 1$. $E_t = E + E = 2/\sqrt{(\Delta\varepsilon/(1-\varepsilon))} = W^+$ mass. $E_i = E - E$ gives E&M that also interacts weakly with weak force.

IIIrd →IV quadrant rotation is the W^- . Do the math and get a Proca equation again.

$E = 1/\sqrt{\kappa_{00}} - 1 = [1/\sqrt{(1-\Delta\varepsilon/(1-\varepsilon)-r_H/r)}] - 1 = [1/\sqrt{(\Delta\varepsilon/(1-\varepsilon))}] - 1$. $E_t = E + E = 2/\sqrt{(\Delta\varepsilon/(1-\varepsilon))} = W^-$ mass. $E_i = E - E$ gives E&M that also interacts weakly with weak force.

II → III quadrant rotation is the Z_0 . Do the math and get a Proca equation. C_M charge cancelation. D14 gives $1/(1+\varepsilon)$ gives 0 charge since $\varepsilon \rightarrow 1$ to case 1 in appendix C2.

$E = 1/\sqrt{\kappa_{00}} - 1 = [1/\sqrt{(1-\Delta\varepsilon/(1+\varepsilon)-r_H/r)}] - 1 = [1/\sqrt{(\Delta\varepsilon/(1+\varepsilon))}] - 1$. $E_t = E + E = 2/\sqrt{(\Delta\varepsilon/(1+\varepsilon))} - 1 = Z_0$ mass. $E_i = E - E$ gives E&M that also interacts weakly with weak force. Seen in small left handed polarization rotation of light.

IV→I quadrant rotation through those 2 neutrinos gives 2 objects. $r_H=0$

$E = 1/\sqrt{\kappa_{00}} - 1 = [1/\sqrt{(1-\Delta\varepsilon/(1+\varepsilon))}] - 1 = \Delta\varepsilon/(1+\varepsilon)$. Because of the +- square root $E = E + -E$ so E rest mass is 0 or $\Delta\varepsilon = (2\Delta\varepsilon)/2$ reduced mass.

$E_t = E + E = 2E = 2\Delta\varepsilon$ is the pairing interaction of SC. The $E_i = E - E = 0$ is the 0 rest mass photon Boson. Do the math (eq.A2-A7) and get Maxwell's equations. Note there was no charge C_M on the two ν s. Note we get SM particles out of composite e,ν using required eq.9 rotations for

A6 Object B Effect On Inertial Frame Dragging (from appendix D)

The fractal implications are that we are inside a cosmological positron inside a proton $2P_{3/2}$ at $r=r_H$ state. The cosmological object (electron) we are inside of is a positron and call it object A which orbits electron object B with a given distant 3rd object C. Object B is responsible for the mass of the electron since it's frame dragging creates that Kerr metric $(a/r)^2 = m_e c^2$ (D9) result used in eq.D9. So Newpde ground state $m_e c^2 \equiv \langle H_e \rangle$ is the fundamental Hamiltonian eigenvalue defining idea for composite e,ν, $r=r_H$ implying Fermi 4 point $E = \int \psi^\dagger H \psi dV = \int \psi^\dagger \psi H dV = \int \psi^\dagger \psi G$

Recall for composite e,ν all interactions occur inside r_H $(4\pi/3)\lambda^3 = V_{rH}$. $\frac{1}{v^{1/2}} = \psi_e = \psi_3 \frac{1}{v^{1/2}} =$

$$\psi_\nu = \psi_4 \text{ so 4pt } \iiint_0^{r_H} \psi_1 \psi_2 \psi_3 \psi_4 dV = 2G \iiint_0^{r_H} \psi_1 \psi_2 \frac{1}{v^{1/2}} \frac{1}{v^{1/2}} V$$

$$\equiv \iiint_0^{r_H} \psi_1 \psi_2 G \equiv \iiint_0^{r_H} \psi_1 \psi_2 (2m_e c^2) dV_{rH} = \iiint_0^{r_H} \psi_1 (2m_e c^2) \psi_2 dV_{rH} \quad (A8)$$

Object C adds its own spin (eg., as in 2nd derivative eq.A1) to the electron spin (1,IV quadrants) and the W associated with the 2P_{3/2} state at r=r_H thereby adds a derivative in a neutrino quadrant (fig.4) thereby including neutrinos in the Fermi 4pt. So 2nd derivative

$$\Sigma((\gamma^\mu \sqrt{\kappa_{\mu\mu}} dx_\mu) - i\kappa)(\gamma^\nu \sqrt{\kappa_{\nu\nu}} dx_\nu + i\kappa)\chi = \Sigma((\gamma^\mu \sqrt{\kappa_{\mu\mu}} dx_\mu) - i\kappa)\psi \text{ so } \frac{1}{2}(1 \pm \gamma^5)\psi = \chi. \quad (A9)$$


In that regard the expectation value of γ^5 is speed and varies with $e^{i3\phi/2}$ in the trifoldium. The spin $\frac{1}{2}$ decay proton $S_{1/2} \propto e^{i\phi/2} \equiv \psi_1$, the original ortho 2P $\frac{1}{2}$ particle is chiral $\chi = \psi_2 \equiv \frac{1}{2}(1 - \gamma^5)\psi = \frac{1}{2}(1 - \gamma^5 e^{i3\phi/2})\psi$. Initial 2P $\frac{1}{2}$ electron ψ is constant. Start with initial ortho state χ . These γ^5 terms then modify equation A8 to read

$$= \iiint_0^{V_{rH}} \psi_1 \psi_2 (2m_e c^2) dV_{rH} = \iint \psi_{S1/2}^* (2m_e c^2 V_{rH}) \chi dV_\phi = K \int \langle e^{i\frac{\phi}{2}} [\Delta \epsilon V_{rH}] (1 - \gamma^5 e^{i\frac{\phi}{2}}) \psi \rangle d\phi = K G_F \int \langle e^{i\phi/2} - \gamma^5 i e^{i(4/2)\phi} \rangle d\phi = K G_F \left(\frac{2e^{i\phi}}{i} \Big|_0^{2\pi+c} - \frac{2\gamma^5 e^{i4\phi}}{i4} \Big|_0^{2\pi+c} \right) = k(1/4 + i\gamma^5) = k(.225 + i\gamma^5 0.974) = k(\cos 13^\circ + i\gamma^5 \sin 13^\circ) \text{ deriving the } 13^\circ \text{ Cabbibo angle.}$$

With previously mentioned CP result (direct evidence of fractal universe) get CKM matrix

A7 Object C Effect on Inertial Frame Dragging and G_F found by using eq.A8 again (N=1 ambient cosmological metric)

Review of 2P_{3/2} Next higher fractal scale (X10⁴⁰), cosmological scale. Recall from D9 $m_e c^2 = \Delta \epsilon$ is the energy gap for object B vibrational stable eigenstates of composite 3e (vibrational perturbation r is the only variable in Frobenius solution, part II Ch.8,9,10) proton. Observer in object A. $\Delta m_e c^2$ gap = object C scissors eigenstates. is what we see at object A but $\Delta m_e c^2$ gets boosted by γ by rotation into the object B direction. (to compare with the object B $m_e c^2$ gap).



From fig 7 $r^2 = 1^2 + 1^2 + 2(1)(1)\cos 120^\circ = 3$, so $r = \sqrt{3}$. Recall for the positron motion $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = 917$.

So start with the distances we observe which are the Fitzgerald contracted $AC =$

$$r_{CA} = 1 \sqrt{1 - \frac{\cos^2 30^\circ c^2}{c^2}} \sqrt{3} = .866 = \cos 30^\circ = CA \text{ and Fitzgerald contracted } AB = r_{BA} = x/\gamma = 1/\gamma \text{ so for}$$

Fitzgerald contracted $x=1$ for AB (fig7). We can start at $t=0$ with the usual Lorentz transformation for the time component.

$$t' = \gamma(ct - \beta x) = kmc^2.$$

since time components are Lorentz contracted proportionally also to mc^2 , both with the γ multiplication.

In the object A frame of reference we see $\Delta m_e c^2$ which is the average of left and right object C motion effect. We go into the AB frame of reference to compare the object B $m_e c^2$ with this $\Delta m_e c^2$. Going into the AB frame automatically boosts $\Delta m_e c^2$ to $\gamma \Delta m_e c^2$. So start from a already Fitzgerald contracted x/γ . Next do the time contraction γ to that frame:

$$t'' = k\gamma \Delta m_e c^2 = \gamma \beta r_{AB} = \gamma \beta \left(\frac{x}{\gamma} \right) = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \beta \left(\sqrt{1 - \frac{v^2}{c^2}} \sqrt{1} \right) = \beta$$

with k defining the projection of tiny $\Delta m_e c^2$ “time” CA onto BA = $\cos\theta$ = projection of BA onto CA. But $m_e c^2$ is the result of object B of both of the motion and inertial frame dragging reduction (D9) so its γ is large. To make a comparison of ΔE to AB mass $m_e c^2$ CA is rotated and translated to the high speed AB direction and distance with its large γ so thereby *object C becomes mathematically object B* with the same k because of these projection properties of: CA onto BA. So we define projection k from projection of $m_e c^2$: So again

$$t' = \gamma(ct - \beta x) = k m_e c^2 = t' = k m_e c^2 = \gamma \beta r_{CA} = \left(\frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \right) \beta \left(\sqrt{1 - \frac{\cos^2 30^\circ c^2}{c^2}} \sqrt{3} \right) = \gamma \beta \cos 30^\circ$$

Take the ratio of $\frac{k \gamma \Delta m_e c^2}{k m_e c^2}$ to eliminate k: thus

$$\frac{k \gamma \Delta m_e c^2}{k m_e c^2} = \frac{\gamma \beta \left(\frac{x}{\gamma} \right)}{\gamma \beta r_{CA}} = \frac{1 \beta 1}{\gamma \beta \cos 30^\circ} = \frac{1}{\gamma \cos 30^\circ} \quad \text{so}$$

$$\Delta m_e c^2 = \frac{\beta m_e c^2}{\beta \cos 30^\circ \gamma^2} = \frac{\left(1 - \frac{v^2}{c^2} \right) m_e c^2}{\cos 30^\circ} \quad \text{(A10)}$$

allowing us to finally compare the energy gap caused by object C ($\Delta m_e c^2$) to the energy gap caused by object B ($m_e c^2$. A8). So to summarize $\Delta E = (m_e c^2 / ((\cos 30^\circ)^2 917^2)) = m_e c^2 / 728000$. So the energy gap caused by object C is $\Delta E = (m_e c^2 / ((\cos 30^\circ)^2 917^2)) = m_e c^2 / 728000$. The weak interaction thereby provides the ΔE perturbation ($\int \psi^* \Delta E \psi dV$) inside of r_H creating those Frobenius series (partII) $r \neq 0$ states, for example in the unstable equilibrium $2P_{1/2}$ electrons m_e . so in the context of those e, v rotations giving W and Z_0 . The G can be written for E&M decay as $(2m_e c^2) X V_{TH} = 2m_e c^2 [(4/3)\pi r_H^3]$. But because this added object C rotational motion is eq.A9 Fermi 4 point it is entirely different than a mere ‘weak’ E&M. So for weak decay from equation A8 it is $G_F = (2m_e c^2 / 728,000) V_{TH} = G_F = 1.4 \times 10^{-62} \text{ J-m}^3 = 9 \times 10^{-4} \text{ MeV-F}^3$ **the strength of the Fermi 4pt weak interaction constant** which is the coupling constant for the Fermi 4 point weak interaction integral. Note $2m_e c^2 / 729,000 = 1.19 \times 10^{-19} \text{ J}$. So $\Delta E = 1.19 \times 10^{-19} / 1.6 \times 10^{-19} = .7 \text{ eV}$ which is our ΔE gap for the weak interaction inside the Fermi 4pt. integral for G_F . This ΔE generates that r perturbation (instability) states in the Frobenius solution (partII) and so weak decay. interaction integral. Note $2m_e c^2 / 729,000 = 1.19 \times 10^{-19} \text{ J}$. So $\Delta E = 1.19 \times 10^{-19} / 1.6 \times 10^{-19} = .7 \text{ eV}$ which is our ΔE gap for the weak interaction inside the Fermi 4pt. integral for G_F .

The perturbation r in the Frobenius solution is caused by this ΔE in ($\int \psi^* \Delta E \psi dV$) with available phase space for $\psi^* = \psi_p \psi_e \psi_\nu$. and $\psi = \psi_N$.

A8 Derivation of the Standard Model from Newpde but with No Free parameters

Since we have now derived M_W, M_Z and their associated Proca equations, and Dirac equations for m_τ, m_μ, m_e etc., and G, G_F, k_e^2 Maxwell’s equations, etc. we can now write down the usual Lagrangian densities that implies these results. In the formulation $M_Z = M_W / \cos\theta_w$ you can find the Weinberg angle $\theta_w, g \sin\theta_w = e, g' \cos\theta_w = e$; solve for g and g' , etc., We will have thereby derived the standard model from first principles (i.e., postulate 1). **It no longer contains free parameters.**

Note $C_M = \text{Feynman pt}$ really is the $U(1)$ charge and equation 12 rotation is on the complex plane so it really implies $SU(2)$ (A1) with the sect.3.2 2D eqs. 7+8 = $G_{00} = E_c + \sigma \cdot p_f = 0$ gets the left handedness. Recall the genius of the SM is getting all those properties (of χ, Z_0, W^+, W^-) from

SU(2)XU(1)_L so we really have completely derived the electoweak standard model from eq.12 which comes out of the Newpde given we even found the magnitude of its input parameters (eg., G_F (appendix A7), Cabbibo angle A6).

Appendix B ultimate Occam's razor (observable) also implies the underlying relational N=0 postulate 1 (observable) can also be used in a list-define math to get the *real number* algebra (without all those many Rel#math axioms). Eg., $1 \cup 1 \equiv 1+1$ (Ch.2).

Postulate 1 (observable) so *observer* C so $1 \cup C \equiv 1+C$. with algebraic definition of $1 z=zz$ having both 1,0 as solutions so defining negation \sim with $0=1-1$ Thus we can define

$\sim((A \cup B) \sim B \sim A) \equiv A \cap B$. So we have defined intersection \cap so we have derived set theory.

So in postulate 1 $z=zz$ why did 0 come along for the ride? There is a deeper reason in set theory.

Note \emptyset and 0 aren't really new postulates since they postulate literally "nothing".

Recall we just derived set theory from the postulate of 1 (observable).

The null set \emptyset is the subset of every set. In the more fundamental set theory formulation

$\{\emptyset\} \subset \{\text{all sets}\} \Leftrightarrow \{0\} \subset \{1\}$ since $\emptyset = \emptyset \cup \emptyset \Leftrightarrow 0+0=0$, $\{\{1\} \cup \emptyset\} = \{1\} \Leftrightarrow 1+0=1$.

So list $1 \cup 1 \equiv 1+1 \equiv 2$, $2 \cup 1 \equiv 1+2 \equiv 3$,...all the way up to 10^{82} (see Fiegenbaum point) and **define** all this list as $a+b=c$, etc., to create our algebra and numbers which we use to write [equation 1](#) $z=zz+C$, $\delta C=0$ for example. Recall every set has the null set as a subset.

B2 2D+2D→4D

Note adding the N=0 fractal scale 2D δz perturbation to N=1 eq.7 2D gives curved space 4D. So

$(dx_1+idx_2)+(dx_3+idx_4) \equiv dr+idt$ given (eqs5,7a) $dr^2-dt^2=(\gamma^r dr+i\gamma^t dt)^2$ if $dr^2 \equiv dx^2+dy^2+dz^2$ (3D

orthogonality) so that $\gamma^r dr \equiv \gamma^x dx + \gamma^y dy + \gamma^z dz$, $\gamma^i \gamma^j + \gamma^j \gamma^i = 0$, $i \neq j$, $(\gamma^i)^2 = 1$, rewritten (with curved space $\kappa_{\mu\nu}$ eq.13-15) $(\gamma^x \sqrt{\kappa_{xx}} dx + \gamma^y \sqrt{\kappa_{yy}} dy + \gamma^z \sqrt{\kappa_{zz}} dz + \gamma^t \sqrt{\kappa_{tt}} dt)^2 = \kappa_{xx} dx^2 + \kappa_{yy} dy^2 + \kappa_{zz} dz^2 - \kappa_{tt} dt^2 = ds^2$.

More fundamentally satisfying this 4D Clifford algebra and complex orthogonalization

requirement is a special case of any 2 $x_i x_j$ in eq.3 (directly from postulate1): Imposing

orthogonality thereby creates 6 pairs of eqs.3&5. So each particle carries around it's own $dr+idt$

complex coordinates with them on their world lines. Alternatively this 2D $dr+idt$ is a 'hologram'

'illuminated' by a modulated $dr^2+dt^2=ds^2$ 'circle' wave (as 2nd derivative wave equation

operators from eq.11 circle) since 4Degrees of freedom are imbedded on a 2D (dr,dt) surface

here, with observed coherent superposition output as eq.16 solutions. A more direct way is to

simply write the 4Degrees of freedom on the 2D surface as $dr+idt = (dr_1+idt_1)+(dr_2+idt_2)$

$= (dr_1, \omega dt_2), (dr_2, idt_2) = (x, z, y, idt) = (x, y, z, idt)$, where $\omega dt \equiv dz$ is the z direction spin $\frac{1}{2}$ component ω

(angular velocity) axial vector of the Newpde lepton (eqs.7-9); which we get anyway from lepton equation eq.16.

N=-1 and dimensionality

Note the N=-1 (GR) is yet another δz perturbation of N=0 $\delta z'$ perturbation of N=1 observer

thereby adding at least 1 independent parameter dimension to our $\delta z + (dx_1+idx_2) + (dx_3+idx_4)$

$(4+1)$ explaining why Kaluza Klein 5D $R_{ij}=0$ works so well: GR is really 5D if E&M

included. Note these fractal N=-1 fractal scale wound up balls at $r_H=10^{-58}m$ are a lot smaller than

the Planck length. But if only N=1 observer and N=-1 are used (no N=0) we still have the usual

4D.

Appendix C

Quantum Mechanics Is The Newpde $\psi \equiv \delta z$ (for each N fractal scale)

The postulate of 1 is the source of other properties of $\delta z = \psi$ in addition to those provided

by just the Newpde. For example recall the solution to (postulate 1) $z=zz$ is 1,0. In $z=1-\delta z$,

$\delta z * \delta z$ is (defined as) the probability of z being 0. Recall $z=0$ is the $\xi_0=m_e$ solution(12b) to the

new pde so $\delta z^* \delta z$ is the probability we have just an electron (11b,11c). Note $z=zz$ also thereby conveniently provides us with an automatic normalization of δz . Note also that $(\delta z^* \delta z)/dr$ is also then a one dimensional probability ‘density’. So Bohr’s probability density “postulate” for $\psi^* \psi$ ($\equiv (\delta z^* \delta z)$) is derived here. It is not a postulate anymore. (So Bohr was very close to the postulate of 1, and so using $z=zz$ here.)

Note the electron-positron eq.7 has *two* components (i.e., $dr+dt$ & $dr-dt$,) that both solve eq.5 (and therefore eq.3) *together* as in the $\delta z \equiv \psi = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ singlet state relation with spin S of two electrons $(S_1+S_2)^2 = S^2$. This singlet ψ can be used as a paradigm-model of the iconic idler-signal (Alice and Bob) singlet QM $\delta(p_A-p_B)$ conservation law state, in the Bell’s inequality formulation.. We could then label these two parts of eq.7 *observer* and *object* with associated eq.7 wavefunctions ψ_1, ψ_2 and singlet ψ . Thus we observe ψ_1 (signal) and so infer that there is both ψ_2 (idler from eq.7) and so our singlet wavefunction ψ . So we ‘collapsed’ our wavefunction to ψ by observing it. Then apply the same mathematical reasoning to every other such analog $\delta z \equiv \psi = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ singlet cases (eg., H, V polarized photon emission) and we will also have thereby derived Bell’s inequalities. This is then a derivation of the wave function collapse part of the Copenhagen interpretation of Quantum Mechanics from eq.7 and so from the first principles **postulate 1**.

But this (Copenhagen interpretation) wave function collapse is actually a trivial principle (i.e., so it could be the wave function ψ is trivially just what you measure) except, as EPR pointed out, in this kind of conservation law singlet case laboratory initialization paradigm ψ . To actually know the initial S_1+S_2 in this $\delta z \equiv \psi = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ QM singlet state is actually a **rare (laboratory setting) case** and so its spooky superluminal collapse is not a universal attribute (that being the new fad taking over theoretical physics) of all observed particles. So even the core Bertlmann’s socks situation is rare and without it Bell’s inequalities don’t apply and so in that case there is no such spookiness.

Also recall from appendix A dr^2+dt^2 is a second derivative *operator* wave equation (A1,eq.11) that holds all the way around the circle (even for the eq.10 vacuum solutions), gives waves. In eq.12, error magnitude C (sect.2.3) is also a $\delta z'$ angle measure on the dr, idt plane. One extremum ds ($z=0$) is at 45° so the largest C is on the diagonals (45°) where we have eq.5 extremum holding: particles. So a wide slit has high uncertainty, so large C (rotation angle) so we are at 45° (eg., particles, eq.16 photoelectric effect). For a *small slit* we have less uncertainty so smaller C, not large enough for 45° , so only the *wave equation* A1 holds (small slit diffraction). Thus we derived wave particle duality here. So complementarity is derived here, not postulated. Recall wave equation eq.A1 iteration of the New pde with eq.11 operator formalism. So $dr/ds=k$ in the sect.1 $\delta z = dse^{i\theta}$ exponent then becomes $k=2\pi/\lambda$. Multiplying both sides by \hbar with $\hbar k = mv$ as before we then have the DeBroglie equation that relates particle momentum to wavelength in quantum mechanics. Equation 8a (sect.1) then counts units N of $(dt/ds) = \hbar\omega = \hbar ck$ on the diagonal so that $E = p v = \hbar\omega$ for all energy components, universally. Thus this eq.11a counting N does not require the (well known) quantization of the E&M field with SHM. First, set the unit of distance r_H on our baseline fractal scale: (eq.1 $N=0$. See figure 1 attachment.). The 4X Mandelbrot set formulation allows only these finite extremum.

$\delta z \equiv \psi$

Appendix D. N=1 observer (eq.13,14,15 give our **Newpde metric** $\kappa_{\mu\nu}$ at $r < r_H, r > r_H$)

Found GR from eq.13 and eq.14 so we can now write the Ricci tensor R_{uv} (and self similar perturbation Kerr metric since frame dragging decreased by external object B, sect.A6). Also for fractal scale $N=0$, $r_H=2e^2/m_e c^2$, and for $N=-1$ $r'_H=2Gm_e/c^2=10^{-40}r_H$.

Nonzero Generic maximally symmetric (MS) ambient metric (meaning $N=1$) generated by object B

$N=2$ big guy sees us from the outside and so sees a sine oscillation eq.17. To see what we see($N=1$) he multiplies \sin by i and u by ' i ' since we are inside (so since in eq. 17->17a then $i \sin u \rightarrow \sin iu$). So start simple with complete frame dragging suppression eq.13, 15 but with ambient metric (provided by later **perturbation** $a \ll r$ **provided by some rotation**) metric ansatz: $ds^2 = -e^\lambda (dr)^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2 + e^\mu dt^2$ so that $g_{00}=e^\mu$, $g_{rr}=e^\lambda$. From eq. $R_{ij}=0$ for spherical symmetry in free space

$$R_{11} = \frac{1}{2}\mu'' - \frac{1}{4}\lambda' \mu' + \frac{1}{4}(\mu')^2 - \lambda'/r = 0 \quad (D1)$$

$$R_{22} = e^{-\lambda} [1 + \frac{1}{2} r(\mu' - \lambda')] - 1 = 0 \quad (D2)$$

$$R_{33} = \sin^2 \theta \{ e^{-\lambda} [1 + \frac{1}{2} r(\mu' - \lambda')] - 1 \} = 0 \quad (D3)$$

$$R_{00} = e^{\mu-\lambda} [-\frac{1}{2}\mu'' + \frac{1}{4} \lambda' \mu' - \frac{1}{4}(\mu')^2 - \mu'/r] = 0 \quad (D4)$$

$$R_{ij} = 0 \text{ if } i \neq j$$

(eq. D1 -D4 from pp.303 Sokolnikof(8)): Equation D2 is a mere repetition of equation D3. We thus have only three equations on λ and μ to consider. From equations D1, D4 we deduce that $\lambda' = -\mu'$ so that radial $\lambda = -\mu + \text{constant} = -\mu + C$ where C represents a possible \sim constant ambient metric contribution which could be imaginary in the case of the slowly oscillating ambient metric of nearby object B. So $e^{-\mu+C} = e^\lambda$. Then D2 can be written as:

$$e^{-C} e^\mu (1 + r\mu') = 1 \quad (D5)$$

Set $e^\mu = \gamma$. So $e^{-\lambda} = \gamma e^{-C}$ and so integrating this first order equation (equation.D11) we get:

$$\gamma = -2m/r + e^C \equiv e^\mu = g_{00} \text{ and } e^{-\lambda} = (-2m/r + e^C) e^{-C} = 1/g_{rr} \quad (D6)$$

From equation D6 we can identify radial C with also rotational Kerr metric oblateness perturbation Mandelbulb component here (D8 below) of Mandelbrot set Fig.6 eq.18 $2m/r = r_H/r = C_M/\xi r = e^{-C} = e^{-(\varepsilon+\Delta\varepsilon)} = \tau + \mu + \Delta\varepsilon$. (eq.17a). We end up being at the horizon r_H in equation D8. So $2m/r$ is set equal to e^C in eq. D6. So at the end, at the horizon r_H , in eq.D8, $2m/r$ is set equal to $e^C = e^{-(\varepsilon+\Delta\varepsilon)}$ in D6. So $\kappa_{00} = 1 - e^{-(\varepsilon+\Delta\varepsilon)} - 2m/r$. from eq.17. Given external object B oscillating zitterbewegung for $r < r_C$ then $e^{-(\varepsilon+\Delta\varepsilon)} \rightarrow e^{-i(\varepsilon+\Delta\varepsilon)}$ so that $\kappa_{00} = 1 - e^{-i(\varepsilon+\Delta\varepsilon)} - 2m/r$ (D7) So: $e^{-\lambda} = 1/\kappa_{rr} = 1/(1-2m'/r)$

Perturbative self similar rotation providing the above ambient metric Generated by object B $N=1$ observer scale

Our new pde has spin S and so the self similar ambient metric on the $N=0$ th fractal scale is the Kerr metric which contains those ambient metric **perturbation rotations** ($d\theta/dt$ T violation so (given CPT) then **CP violation**)

$$ds^2 = \rho^2 \left(\frac{dr^2}{\Delta} + d\theta^2 \right) + (r^2 + a^2) \sin^2 \theta d\phi^2 - c^2 dt^2 + \frac{2mr}{\rho^2} (a \sin^2 \theta d\theta - c dt)^2, \quad (D8)$$

where $\rho^2(r, \theta) \equiv r^2 + a^2 \cos^2 \theta$; $\Delta(r) \equiv r^2 - 2mr + a^2$, In our 2D $d\phi=0$, $d\theta=0$ Define:

$$\left(\frac{r^2 + a^2 \cos^2 \theta}{r^2 - 2mr + a^2} \right) dr^2 + \left(1 - \frac{2m}{r^2 + a^2 \cos^2 \theta} \right) dt^2 \quad \theta \neq 0$$

$\rho^2(r, \theta) \equiv r^2 + a^2 \cos^2 \theta$; $\Delta(r) \equiv r^2 - 2mr + a^2$, $r^{\wedge 2} \equiv r^2 + a^2 \cos^2 \theta$, $r'^2 \equiv r^2 + a^2$. Inside r_H $a \ll r$, $r \gg 2m$

$$\left(\frac{(r')^2}{(r'r)^2-2mr}\right) dr^2 + \left(1 - \frac{2mr}{(r')^2}\right) dt^2 + \dots = \left(\frac{1}{\frac{(r')^2}{(r')^2} \frac{2mr}{(r')^2}}\right) dr^2 + \left(1 - \frac{2mr}{(r')^2}\right) dt^2.$$

The $(r'/r')^2$ term is

$$\frac{(r')^2}{(r')^2} = \frac{r^2+a^2}{r^2+a^2\cos^2\theta} = \frac{1+\frac{a^2}{r^2}}{1+\frac{a^2}{r^2}\cos^2\theta} \approx 1/g_{rr}(\approx g_{00}) \text{ From D7: } \xi_1 = e^{i(\varepsilon+\Delta\varepsilon)} \text{ for } e^C = e^{i(\varepsilon+\Delta\varepsilon)}$$

= $\tau+\mu+\Delta\varepsilon$ =zitterbewegung from D6. $2m/r+e^C$

$$\left(1 + \frac{a^2}{r^2}\right) \left(1 - \frac{a^2}{r^2}\cos^2\theta\right) + \dots = 1 - \frac{a^4}{r^4}\cos^2\theta - \frac{a^2}{r^2}\cos^2\theta + \frac{a^2}{r^2} + \dots = 1 + \frac{a^2}{r^2}(1 - \cos^2\theta) + \dots$$

$$= 1 + \frac{a^2}{r^2}\sin^2\theta + \dots \equiv 1 + \frac{a^2}{r^2}u^2 = (D7,17) = 1 + e^C = 1 + e^{i(\varepsilon+\Delta\varepsilon)} =$$

(Replace a^2/r^2 Kerr object B term with inertial frame D7 dragging mass ξ_1 . In eq.D8 subtract $2mr/(r')^2=r_H/r_H$). In From eq.17a general the closer object B is the larger e^C is.

$$= 1 + \xi_1 - \frac{r_H}{r_H} = e^C = 1 + \varepsilon + \Delta\varepsilon + \dots = e^{i(\varepsilon+\Delta\varepsilon)} \quad (D9)$$

So this is a Kerr metric inertial frame dragging suppression due to outside object B of magnitude $((a/r)\sin\theta)^2 = 1/g_{rr}=e^{i\varepsilon}$ from D7 in the proper frame. In the $N=1$ observer scale at $r=r_H$. Inside object A. ε also changes with time (Mercuron equation D15).

Object B oscillation sound wave observed compression in Shapely, Bootes, rarefaction in Eridanis.

D2 Examples of this ambient metric. N=0 Composite 3e

Introduction: $N=0$ Frobenius solution is for constant ψ (and so constant ε) just inside r_H .

Equations D6,D7,D9 provide the $e^{i(\varepsilon+\Delta\varepsilon)}$ contributions from each maximal symmetry ε source, with the B flux quantization causing the $n\varepsilon$ quantization of the ambient metric. There appear to be 2 B field sources, the two fast moving positrons (are right on r_H and so are close to these boundaries) creating that huge internal magnetic field. So for the inside $1+2(\varepsilon+\Delta\varepsilon)$ get added and we normalize the maximal symmetry B field away for the observer 2nd positron by dividing by $1+\varepsilon$.

In contrast for just *outside* r_H the flux is canceled out because of the frequent creation and annihilation events inside resulting in a Faraday's law B flux change cancellation application that gives the Meisner effect zero point energy (eq.9.22) pion ε' cloud who's energy is thereby added to $2m/r=r_H/r$ as implied by eq. D6. Thus:

For $z=0$ *just inside* r_H , the *two* positrons *each* have constant ψ ($N=0$ ch.8,9) inside r_H . So from eq.D9 divide κ_{rr} by $1+\varepsilon+\varepsilon=1+2\varepsilon.=e^C$ So $\frac{1}{\kappa_{rr}} = (1)(1+2\varepsilon) \equiv 1+2(\varepsilon+\Delta\varepsilon)$ (D9a)

Note negative potential energy here. Normalize out the κ_{00} magnetic field maximal symmetry of the observer by multiplying κ_{00} by $1+\varepsilon=e^C$ for the magnetic (see partII flux of B)

$$\frac{1}{\left(\frac{1+2\varepsilon+\Delta\varepsilon}{1+\varepsilon}-2m/\xi_0 r\right)} dr^2 + (1 - 2m/r\xi_0) dt^2 = \frac{1}{\left(1+\frac{\varepsilon}{1+\varepsilon}-2m/\xi_0 r\right)} dr^2 + \left(1 - \frac{2m}{r\xi_0}\right) dt^2$$

$$= \frac{1}{(1+\varepsilon)-2m/\xi_0 r} dr^2 + \left(1 - \frac{2m}{r\xi_0}\right) dt^2, \quad \varepsilon' \equiv \varepsilon/(1+\varepsilon). \quad (D10)$$

For $z=0$ *just outside* r_H , Since randomly the B field disappears ($dB/dt \neq 0$) due to that creation-annihilation we have a Faraday's law Meisner effect. With outside r_H B results, just divide by $1+\varepsilon'$ (D9) for zero point energy $\varepsilon'' = .08 \pi^\pm$ of eq.9.22 (partII) which has to itself increase and

decrease with (see D9) each of these annihilation events and π^\pm exists just outside r_H (from our Frobenius solution): $\frac{1}{(1+\varepsilon''-2m/\xi_0 r)} dr^2 + ((1 - 2m/\xi_0 r)) dt^2 = ds^2$ (D11)

For $z=0 \rightarrow z=1$ $r \gg r_H$ then free space boost sect.2 $\xi_0 \rightarrow \tau$. Define $\varepsilon' \equiv \frac{\varepsilon}{1+\varepsilon}$. Must normalize again (for local ambient metric $\Delta\varepsilon$ change contributions) so multiply by $\frac{1}{1+\varepsilon'}$ (see D9 for $z=1$ outside)

$$\frac{1}{(1+\frac{\Delta\varepsilon}{1+\varepsilon}-2m/\xi_1 r)} dr^2 + (1 - 2m/r\xi_1) dt^2 = \frac{1}{(1+\frac{\Delta\varepsilon}{1+\varepsilon}-2m/\xi_1 r)} dr^2 + \left(1 - \frac{2m}{r\xi_1}\right) dt^2 \quad (D12)$$

D3 A N=0 Application example: (mentioned on first page)

Separation Of Variables On New Pde

After separation of variables the “r” component of equation 16 (Newpde) can be written as:

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] F - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{j+3/2}{r} \right) f = 0 \quad D13$$

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] f + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{j-1/2}{r} \right) F = 0. \quad D14$$

Using the above Dirac equation component we find the anomalous gyromagnetic ratio Δgy for the spin polarized $F=0$ case. Recall the usual calculation of rate of the change of spin S gives $dS/dt \propto m \propto gyJ$ from the Heisenberg equations of motion. We note that $1/\sqrt{\kappa_{rr}}$ rescales dr in $\left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{j+3/2}{r} \right) f$ in equation C5. Thus to have the same rescaling of r in the second term we must multiply the second term denominator (i.e., r) and numerator (i.e., $J+3/2$) each by $1/\sqrt{\kappa_{rr}}$ and set the numerator ansatz equal to $(j+3/2)/\sqrt{\kappa_{rr}} \equiv 3/2 + J(gy)$, where gy is now the gyromagnetic ratio. This makes our equation D13, D14 compatible with the standard Dirac equation allowing us to substitute the gy into the Heisenberg equations of motion for spin S : $dS/dt \propto m \propto gyJ$ to find the correction to dS/dt . Thus again:

$$\begin{aligned} [1/\sqrt{\kappa_{rr}}](3/2 + J) &= 3/2 + Jgy, \text{ Therefore for } J = 1/2 \text{ we have:} \\ [1/\sqrt{\kappa_{rr}}](3/2 + 1/2) &= 3/2 + 1/2 gy = 3/2 + 1/2(1 + \Delta gy) \end{aligned} \quad D15$$

Then we solve for Δgy and substitute it into the above dS/dt equation.

Thus solve eq. D12, D15 with eq.19 values in $\sqrt{\kappa_{rr}} = 1/\sqrt{(1+\Delta\varepsilon/(1+\varepsilon))} = 1/\sqrt{(1+\Delta\varepsilon/(1+0))} = 1/\sqrt{(1+.0005799/1)}$. Thus from equations C1, D13, D15, A0:

$[\sqrt{(1+.0005799)}](3/2 + 1/2) = 3/2 + 1/2(1 + \Delta gy)$. Solving for Δgy gives anomalous **gyromagnetic ratio correction of the electron** $\Delta gy = .00116$.

If we set $\varepsilon \neq 0$ (so $\Delta\varepsilon/(1+\varepsilon)$) instead of $\Delta\varepsilon$ in the same κ_{00} in eq.16 we get the anomalous **gyromagnetic ratio correction of the muon** in the same way.

Composite 3e: Meisner effect For B just outside r_H . (where the zero point energy particle eq. 9.22 is $.08 = \pi^\pm$) See D11

Composite 3e CASE 1: Plus $+r_H$, therefore is the proton + charge component. Eq. C1 & D11, A0 $1/\kappa_{rr} = 1 + r_H/r_H + \varepsilon'' = 2 + \varepsilon''$. $\varepsilon'' = .08$ (eq.9.22). Thus from eq. C7: $\sqrt{2 + \varepsilon''}(1.5 + .5) = 1.5 + .5(gy)$, $gy = 2.8$

The gyromagnetic ratio of the proton

Composite 3e CASE 2: negative r_H , thus charge cancels, zero charge:

$$\begin{aligned} 1/\kappa_{rr} &= 1 - r_H/r_H + \varepsilon'' = \varepsilon'' \text{ Therefore from equation D15 and case 1 eq.12 } 1/\kappa_{rr} = 1 - r_H/r_H + \varepsilon'' \\ \sqrt{\varepsilon''} (1.5 + .5) &= 1.5 + .5(gy), \text{ } gy = -1.9. \end{aligned}$$

the **gyromagnetic ratio of the neutron** with the other charged and those ortho neutral hyperon magnetic moments scaled using their masses by these values respectively.

D4 Separation of Variables

After separation of variables the “r” component of equation 16 (Newpde) can be written as

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) + m_p \right] F - \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} + \frac{j+3/2}{r} \right) f = 0 \quad D16$$

$$\left[\left(\frac{dt}{ds} \sqrt{\kappa_{00}} m_p \right) - m_p \right] f + \hbar c \left(\sqrt{\kappa_{rr}} \frac{d}{dr} - \frac{j-1/2}{r} \right) F = 0. \quad D17$$

Comparing the flat space-time Dirac equation to the left side terms of equations C5 and C6:

$$(dt/ds)\sqrt{\kappa_{00}}=(1/\kappa_{00})\sqrt{\kappa_{00}}=(1/\sqrt{\kappa_{00}})=\text{Energy}=E \quad D18$$

Note for electron motion around hydrogen proton $mv^2/r=ke^2/r^2$ so $KE=1/2mv^2=(1/2)ke^2/r=PE$

potential energy in $PE+KE=E$. So for the electron (but not the tauon or muon that are not in this

orbit) $PE_e=1/2e^2/r$. Here write the hydrogen energy and pull out the electron contribution. So in eq.B1

and D18 $r_H=(1+1+.5)e^2/(m_\tau+m_\mu+m_e)/2=2.5e^2/(2m_p c^2)$. D19

Variation $\delta(\psi^*\psi)=0$ At $r=n^2a_0$

Next note for the variation in $\psi^*\psi$ is equal to zero at maximum $\psi^*\psi$ probability density where

for the hydrogen atom is at $r=n^2a_0=4a_0$ for $n=2$ and the $\psi_{2,0,0}$ eigenfunction. Also recall eq.B1

eq.19, $\xi_1=m_L c^2=(m_\tau+m_\mu+m_e)c^2=2m_p c^2$ normalizes $1/2ke^2$ (Thus divide $\tau+\mu$ by 2 and then

multiply the whole line by 2 to normalize the $m_e/2$.result. $\varepsilon=0$ since no muon ε here.): Recall in

eq.19 ξ_0 has to be pulled in a Taylor expansion as an operator since it a separate observable. So

substituting eqs.D16,C1 and eq.D12for κ_{00} , and B1,eq19 values in eq.D18:

$$E_e = \frac{(tauon + muon)\left(\frac{1}{2}\right)}{\sqrt{1 - \frac{r_{H'}}{r}}} - (tauon + muon + PE_\tau + PE_\mu - m_e c^2) \frac{1}{2} =$$

$$2(m_\tau c^2 + m_\mu c^2) \frac{1}{2} + 2 \frac{m_e c^2}{2} + 2 \frac{2.5e^2}{2r(m_L c^2)} m_L c^2 - 2 \frac{2e^2}{2r(m_L c^2)} m_L c^2 - 2 \frac{3}{8} \left(\frac{2.5e^2}{r m_L c^2} \right)^2 m_L c^2$$

$$- 2(m_\tau c^2 + m_\mu c^2) \frac{1}{2}$$

$$= \frac{2m_e c^2}{2} + 2 \frac{e^2}{4r} - 2 \frac{3}{8} \left(\frac{2.5}{r m_L c^2} \right)^2 m_L c^2 = m_e c^2 + \frac{e^2}{2r} - 2 \frac{3}{8} \left(\frac{2.5e^2}{r m_L c^2} \right)^2 m_L c^2$$

So: $\Delta E_e = 2 \frac{3}{8} \left(\frac{2.5}{r m_L c^2} \right)^2 m_L c^2 =$ (Third order $\sqrt{\kappa_{\mu\mu}}$ Taylor expansion term)=

$$\Delta E = 2 \frac{3}{8} \left[\frac{2.5(8.89 \times 10^9)(1.602 \times 10^{-19})^2}{(4(.53 \times 10^{-10}))^2((1.67 \times 10^{-27})(3 \times 10^8))^2} \right]^2 (2(1.67 \times 10^{-27})(3 \times 10^8))^2$$

= $hf=6.626 \times 10^{-34} \times 27,360,000$ so that $f=27\text{MHz}$ Lamb shift.

The other 1050Mhz comes from the zitterbewegung cloud.

Note: Need infinities if **flat space Dirac 1928 equation**. For flat space $\partial g_{ik}/\partial x^j=0$ as a limit. Then

must take field $g^{km}=1/0=\infty$ to get finite Christoffel symbol $\Gamma^{m,ij}=(g^{km}/2)(\partial g_{ik}/\partial x^j + \partial g_{jk}/\partial x^i -$

$\partial g_{ij}/\partial x^k)=(1/0)(0)=\text{undefined}$ but still implying *nonzero* acceleration on the left side of the

geodesic equation: $\frac{d^2 x^\mu}{ds^2} = -\Gamma_{\nu\lambda}^\mu \frac{dx^\nu}{ds} \frac{dx^\lambda}{ds}$ So we need infinite fields for flat space. Thus QED

requires (many such) infinities. But we have in general curved space $g_{ij}=\kappa_{ij}$ in the New pde so do

not require that anything be infinite and yet we still obtain for the third order Taylor expansion

term of $\sqrt{\kappa_{\mu\nu}}$ the Lamb shift and anomalous gyromagnetic ratio correction (see above sections

C2,C3).

So renormalization is a perturbative way (given it's flat space Dirac equation and minimal interaction gauge origins) of calculating these (above) same, *NON*perturbative results, it's a perturbative GR theory. But renormalization gives lots of wrong answers too, eg., 10^{96} grams/cm³ vacuum density for starters. (So we drop it here since we don't need it any longer for the high precision QED results.) In contrast note near the end of reference 5 our $G_{00}=0$ for a 2D MS. Thus a vacuum really is a vacuum. Also that large $\xi_1=\tau(1+\varepsilon')$ in r_H in eq.14 is the reason leptons appear point particles (in contrast to the small ξ_0 in the composite 3e baryons).

D5 N=1 internal Observer cosmological physics from Observer at N=2

From Newpde (eg., eq.1.13 Bjorken and Drell) $i\hbar \frac{\partial \psi}{\partial t} = \frac{\hbar c}{i} \left(\alpha_1 \frac{\partial \psi}{\partial x^1} + \alpha_2 \frac{\partial \psi}{\partial x^2} + \alpha_3 \frac{\partial \psi}{\partial x^3} \right) +$

$\beta mc^2 \psi = H\psi$. For electron at rest: $i\hbar \frac{\partial \psi}{\partial t} = \beta mc^2 \psi$ so: $\delta z = \psi_r = w^r(0) e^{-i\varepsilon_r \frac{mc^2}{\hbar} t}$ $\varepsilon_r=+1,$

$r=1,2; \varepsilon_r=-1, r=3,4$): This implies an oscillation frequency of $\omega=mc^2/\hbar$. So the eq.12 the 45° line has this ω oscillation on that δz rotation. The next higher cosmological independent (but still connected by superposition of speeds) fractal scale N=1 the 45° small Mandelbulb chord ε (Fig6)

is now getting smaller with time $t \propto \varepsilon$ as in a separation of variables result: $i\hbar \frac{\partial \psi}{\partial t} =$

$\beta \sum_N (10^{40N} (\omega t)_{\varepsilon+\Delta\varepsilon}) \psi = \beta \sum_N (10^{40N} m_{\varepsilon+\Delta\varepsilon} c^2/\hbar) \psi$ and so for stationary N=1 $\delta z = \sqrt{\kappa_{00}} dt = e^{-i\varepsilon_r \frac{mc^2}{\hbar} t} \rightarrow e^{i(\varepsilon+\Delta\varepsilon)}$ (18)

On our own fractal cosmological scale we are in the expansion stage of one such oscillation.

Recall $N>0 \equiv$ observer. Here we find what that N=2 fractal scale observer sees what we see if $\sin\mu \rightarrow \sinh\mu$ for $r>r_H$ going to $r<r_H$ in $E=1/\sqrt{\kappa_{00}}=1/\sqrt{(1-r_H/r)}$ since the E in $\delta z = e^{iEt} \equiv e^{i\mu}$ and so μ then becomes imaginary. Recall limit R_{ij} as $r \rightarrow 0$ is the source, where gravity creates gravity in the Einstein equations which becomes the modulation of the DeSitter ball. (6.14.2).

$R_{22} = e^{-\lambda} [1 + \frac{1}{2} r(\mu' - v')] - 1$ with $\mu = v$ (spherical symmetry) and $\mu' = -v'$. So as $r \rightarrow 0$, $\text{Im} R_{22} =$

$\text{Im}(e^\mu - 1) = \mu + \dots = \sin\mu = \mu + \dots$ for outside r_H imaginary μ for small r (at the source) so $\sin\mu$ becomes a gravitational source (gravity itself can create gravity as a feedback mechanism). The N=2 observer then multiplies by i iR_{22} , $-\sin\mu$ and μ to get $R_{22} = -\sinh\mu$ to see what the N=2 observer sees that we see inside r_H so:

$R_{22} = e^{-\lambda} [1 + \frac{1}{2} r(\mu' - v')] - 1 = -\sinh v = -(e^v - e^{-v})/2$, $v' = -\mu'$ so

$e^{-\mu} [-r(\mu')] = -\sinh\mu - e^{-\mu} + 1 = -(e^{-\mu} + e^\mu)/2 - e^{-\mu} + 1 = -(e^{-\mu} + e^\mu)/2 + 1 = -\cosh\mu + 1$. So given $v' = -\mu'$

$e^{-v} [-r(\mu')] = 1 - \cosh\mu$. Thus

$e^{-\mu} r(d\mu/dr) = 1 - \cosh\mu$

This can be rewritten as:

$$e^\mu d\mu / (1 - \cosh\mu) = dr/r \quad (D20)$$

The integration is from $\xi_1 = \mu = \varepsilon = 1$ to the present day mass of the muon = .06 (X tauon mass).

Integrating equation B from $\varepsilon = 1$ to the present ε value we then get:

$$\ln(r_{M+1}/r_{bb}) + 2 = [1/(e^\mu - 1) - \ln[e^\mu - 1]]/2 \quad (D21)$$

then $r_{bb} \approx 50 \text{Mkm} \equiv$ mercuron (initial $r=r_H$ each baryon. Big bang 10^{82} baryons sect.2.3). Solve for r_{M+1} , as function of μ . Find present derivative, find du from Hubble constant normalize the number to 13.7 to find total time u . Find we are now at 370by. This long of time explains the cbr thermalization and mature galaxies at dawn (instead of $\sim 200 \text{My}$ after bb, it is 370by). The zitterbewegung (sound wave) of object B creates the condensation (at the Shapely concentration) and rarefaction void in Eridanus: we are astronomically observing here selfsimilar object B sound waves *inside* of a proton.

After a large expansion from r_{bb} our eq.14 eq.15 Schwarzschild finally becomes **Minkowski** $ds^2=-dx_0^2+\sum_{i=1}^n dx_i^2$. The submanifold is $-x_0^2+\sum_{i=1}^n x_i^2=\alpha^2$

In static coordinates r,t : (the **New pde** zitterbewegung **harmonic coordinates** x_i for $r<r_H$)
 $x_0=\sqrt{(\alpha^2-r^2)}\sinh(t/\alpha)$: (sinht is small t limit of equation D15. 5Tyears is the period>>370by)
 $x_1=\sqrt{(\alpha^2-r^2)}\cosh(t/\alpha)$
 $x_i=rz_i$ $2\leq i\leq n$ z_i is the standard imbedding $n-2$ sphere. R^{n-1} which also implies the **De Sitter** metric: $ds^2=-(1-r^2/\alpha^2)dt^2+(1-r^2/\alpha^2)^{-1}dr^2+d\Omega_{n-2}^2$ (D16) **our observed ambient metric.**

D6 Mixed states of $\Delta\varepsilon$ and ε $N=-1$ outside so $1S_{1/2}$ state with r

$\hbar_{N=-1} \Delta x \Delta(m_{N=-1}c) = \hbar/2$. $m_{N=-1} = 10^{-40} m_e$. So $\Delta x = 10^5 LY$ galaxy. $1S_{1/2}$ state may be flattened since such states are stable since $g_{00} = \kappa_{00}$.

From D13 metric source note $\Delta\varepsilon$ and ε operators so $\Delta\varepsilon\varepsilon$ (operating on Newpde ψ_N) is a new state, a “mixed state” that in the next higher scale classical limit then is a grand canonical ensemble with nonzero chemical potential (i.e., a “mixture” of systems). 2nd derivative of $\cos x = -\cos x$ so $\Delta g_{00} = -g_{00} = \cos \Delta\varepsilon$. That $g_{00} = \kappa_{00}$ in the *halo of the Milky Way galaxy* is the fundamental equation of metric quantization for all the multiples of 100 metric quantization, but here for $r < r_H$. So in general $\kappa_{00} = e^{i(m_e + m_\mu)}$, $m_e = .000058$ is the electron mass (as a fraction of the Tauon mass.) which is the component in the resulting m_e, m_μ operator sequence.

Imaginary part R_{22} locally for 2D MS $R_{00} = \Delta g_{00} = \kappa_{00}(R/2) = \cos \Delta\varepsilon$ gives also the local mixed $\Delta\varepsilon, \varepsilon$ states of part III metric quantization. Set $\cos(\Delta\varepsilon/(1-2\varepsilon)) = \kappa_{00} = g_{00}$, $mv^2/r = GMm/r^2$ so $GM/r = v^2$ COM in the galaxy halo (circular orbits) $(1/(1-2\varepsilon))$ term from D9a just inside r_H so

Pure state $\Delta\varepsilon$ (ε excited $1S_{1/2}$ state of ground state $\Delta\varepsilon$, so not same state as $\Delta\varepsilon$)

$Rel \kappa_{00} = \cos \mu$ from D9, A0

$$\text{Case 1 } 1-2GM/(c^2r) = 1-2(v/c)^2 = 1-(\Delta\varepsilon/(1-2\varepsilon))^2/2 \quad (D17)$$

So $1-2(v/c)^2 = 1-(\Delta\varepsilon/(1-2\varepsilon))^2/2$ so $=(\Delta\varepsilon/(1-2\varepsilon))c/2 = .00058/(1-(.06)^2)(3 \times 10^8)/2 = 99 \text{ km/sec} \approx 100 \text{ km/sec}$ (Mixed $\Delta\varepsilon, \varepsilon$, states classically here are grand canonical ensembles with nonzero chemical potential.). For ringed (not hub) galaxies the radial value becomes $100/2 = 50 \text{ km/sec}$.

Mixed state $\varepsilon \Delta\varepsilon$ (Again $GM/r = v^2$ so $2GM/(c^2r) = 2(v/c)^2$.)

$$\text{Case 2 } g_{00} = 1-2GM/(c^2r) = Rel \kappa_{00} = \cos[\Delta\varepsilon + \varepsilon] = 1 - [\Delta\varepsilon + \varepsilon]^2/2 = 1 - [(\Delta\varepsilon + \varepsilon)^2 / (\Delta\varepsilon + \varepsilon)]^2/2 = 1 - [(\Delta\varepsilon^2 + \varepsilon^2 + 2\varepsilon\Delta\varepsilon) / (\Delta\varepsilon + \varepsilon)]^2$$

The $\Delta\varepsilon^2$ is just the above first case (Case 1) so just take the mixed state cross term

$$[\varepsilon\Delta\varepsilon / (\varepsilon + \Delta\varepsilon)] = c[\Delta\varepsilon / (1 + \Delta\varepsilon/\varepsilon)]/2 = c[\Delta\varepsilon + \Delta\varepsilon^2/\varepsilon + \dots \Delta\varepsilon^{N+1}/\varepsilon^{N+1}]/2 = \sum v_N. \text{ Note each term in this expansion is itself a (mixed state) operator. So there can't be a single } v \text{ in the large gradient } 2^{nd} \text{ case so in the equation just above we can take } v_N = [\Delta\varepsilon^{N+1} / (2\varepsilon^N)]c. \quad (D18)$$

From eq. D18 for example $v = m 100^N \text{ km/sec}$. $m=2, N=1$ here (Local arm). In part III we list hundreds of examples of D18: (sun 1,2 km/sec, galaxy halos $m 100 \text{ km/sec}$). The linear mixed state subdivision by this ubiquitous ~ 100 scale change factor in r_{bb} (due to above object B zitterbewegung spherical Bessel function resonance boundary conditions resulting in nodes) created the voids. Same process for $N-1$ (so 100X smaller) antinodes get galaxies, 100X smaller: globular clusters, 100X smaller solar systems, etc., So these smaller objects were also created by mixed state metric quantization (eq. D18) resonance oscillation inside initial radius r_{bb} .

We include the effects of that object B drop in inertial frame dragging on the inertial term m in the Gamow factor and so lower Z nuclear synthesis at earlier epochs ($t > 18$ by)BCE. (see partIII)

Appendix E Δ Modification of Usual Elementary Calculus ϵ, δ ‘tiny’ definition of the limit.

Recall that: given a number $\epsilon > 0$ there exists a number $\delta > 0$ such that for all x in S satisfying

$$|x - x_0| < \delta$$

we have

$$|f(x) - L| < \epsilon$$

Then write $\lim_{x \rightarrow x_0} f(x) = L$

Thus you can take a smaller and smaller ϵ here, so then $f(x)$ gets closer and closer to L even if x never really reaches x_0 . ‘Tiny’ for $h \rightarrow L_1$ and $f(x+h) - f(x) \rightarrow L_2$ then means that $L=0 = L_1$ and L_2 . ‘Tiny’ is this difference limit.

Hausdorf (Fractal) s dimensional measure using ϵ, δ

Diameter of U is defined as $|U| = \sup\{|x - y| : x, y \in U\}$. $E \subset \cup_i U_i$ and $0 < |U_i| \leq \delta$

$$H_\delta^s(E) = \inf \sum_{i=1}^{\infty} |U_i|^s$$

analogous to the elementary $V = U^s$ where of $s=3$, $U=L$ then V is the volume of a cube $\text{Volume} = L^3$. Here however ‘s’ may be noninteger (eg., fractional). The volume here would be the respective Hausdorf outer measure.

The infimum is over all countable δ covers $\{U_i\}$ of E.

To get the Hausdorf outer measure of E we let $\delta \rightarrow 0$ $H^s(E) = \lim_{\delta \rightarrow 0} H_\delta^s(E)$

The restriction of H^s to the σ field of H^s measurable sets is called a Hausdorf s-dimensional measure. Dim E is called the Hausdorf dimension such that

$$H^s(E) = \infty \text{ if } 0 \leq s < \dim E, \quad H^s(E) = 0 \text{ if } \dim E < s < \infty$$

So if s implies a zero H or infinite H it is not the correct dimension. This rule is analogous to the definition of the (fractal) Mandelbrot set itself in which a C that gave infinity is rejected by the definition $\delta C = 0$ we can model as a binary pulse ($z = zz$ solution is binary $z = 1, 0$) with

$zz = z(1)$ is the algebraic definition of 1 and can add real constant C (so $z' = z'z' - C$, $\delta C = 0$ (2)), $z \in \{z'\}$

Plug $z' = 1 + \delta z$ into eq.2 and get $\delta z + \delta z \delta z = C$ (3)

so $\delta z = (-1 \pm \sqrt{1 + 4C}) / 2 = dr + idt$ (4)

for $C < -1/4$ so real line $r=C$ is immersed in the complex plane.

$z = z_0 = 0$ To find C itself substitute z' on left (eq.2) into right $z'z'$ repeatedly & get $z_{N+1} = z_N z_N - C$. $\delta C = 0$ requires us to reject the Cs for which

$-\delta C = \delta(z_{N+1} - z_N z_N) = \delta(\infty - \infty) \neq 0$. **$z = zz$ solution is 1,0** so initial

gets the **Mandelbrot set** C_M (fig2) out to some $\|\Delta\|$ distance from $C=0$. Δ found from $\partial C / \partial t = 0$, $\delta C \equiv \delta C_r = (\partial C_M / \partial (drdt)) dr = 0$ extreme giving the Feigenbaum point $\|C_M\| = \|-1.400115..\|$ global max given this $\|C_M\|$ is biggest of all.

If s is not an integer then the dimensionality it is has a fractal dimension.

But because the Feigenbaum point Δ uncertainty limit is the r_H horizon, which is impenetrable (sect.2.5, partI), ϵ, δ are not dr/ds eq.11a observables for $0 < \epsilon, \delta < r_H$. Instead $\epsilon, \delta > \Delta = r_H$ = the next $10^{40}X$ smaller fractal scale Mandelbrot set at the Feigenbaum point.

Appendix F

Review This is an Occam's razor *optimized* (i.e., $\delta C=0, \|C\|=\text{noise}$)

POSTULATE OF 1

So

$z=zz$ (1) is the algebraic definition of 1, add real constant C (i.e., $z'=z'z', \delta C=0$) (2), $z \in \{z'\}$

Recall from eq.7 that $dr+dt=ds$. So combining in quadrature eqs 7&11 $SNR \delta z = (dr/ds + dt/ds) \delta z = ((dr+dt)/ds) \delta z = (1) \delta z$ (11c, append) and so having come *full circle* back to postulate 1 as a real eigenvalue ($1 \equiv \text{Newpde electron}$). So we really do have a binary physics signal. So, having come *full circle* then: (postulate 1 \Leftrightarrow Newpde)

Digital communication analogy: Binary ($z=zz$) 1,0 signal with white noise $\delta C=0$ in $z'+C=z'z'$. Recall the algebraic definition of 1 is $z=zz$ which has solutions 1,0.(11c). Boolean algebra. Also you could say white noise C has a variation of zero ($\delta C=0$) making it easy to filter out (eg., with a Fourier cutoff filter). So you could easily make the simple digital communication analogy of this being a binary ($z=zz$) 1,0 signal with white noise $\delta C=0$ in $z'+C=z'z'$. (However the noise is added a little differently here ($z+C=zz$) than in statistical mechanics signal theory (eg., There you might use deconvolved signal=convolution integral [(transfer function)signal]dA)). where the 'signal' actually would equal $z+C$, not the usual $(2J_+(r)/r)^2$ psf So this is not quite the same math as in signal theory statistics statistical mechanics.)